

Recombinant Mouse I17RB Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP51922

Size: 10ug, 100ug

Protein Name: Interleukin-17 receptor B (IL-17 receptor B) (IL-17RB) (IL-17 receptor homolog 1) (IL-17ER) (IL-17Rh1) (IL17Rh1) (Interleukin-17B receptor) (IL-17B receptor)

Synonym: Il17rb, Evi27, Il17br

UniProt ID: Q9JIP3

UniProt Link: www.uniprot.org/uniprotkb/Q9JIP3/entry

Species Source: Mouse

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 499 AA

Molecular Weight: 56 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MLLVLLILAASCRSALPREPTIQCGSETGPSPEWMVQHSLTPGDLRDLQVELVKTSVAEEFSILMNI
SWILRADASIRLLKATKICVSGKNNMNSYSCVRCNYTEAFQSQTRPSGGKWTFSYVGFPVELSTLYLI
SAHNIPNANMNEDSPSLSVNFTSPGCLNHVMKYKKQCTEAGSLWDPDITACKKNEKMVEVNFTTNP
LGNRYTILIQRDITLGFSLVLENKLMRTSVAIPVTEESEGAIVQLTPYLHTCGNDCIRREGTVVLCSET
SAPIPPDDNRRMLGGWLPLFLVLLVAVVWLAAGIYLTWRQGRSTKTSFPISMMLPLIKVLVYPSEIC
FHHTVCRFTDFLQNYCRSEVILEKWQKKKIAEMGPVQWLTTQKQAADKVVFLPSDVPTLCDSACG
HNEGSARENSQDLFPLAFNLFCSDFSSQTHLHKYLVVYLGADLKGDYNALSVCPQYHLMKDATAF
HTELLKATQSMSVKKRSQACHDSCSPL

3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.

Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant Interleukin-17 receptor B (IL-17 receptor B) (IL-17RB) (IL-17 receptor homolog 1) (IL-17ER) (IL-17Rh1) (IL17Rh1) (Interleukin-17B receptor) (IL-17B receptor) produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: Receptor for the pro-inflammatory cytokines IL17B and IL17E. May play a role in controlling the growth and/or differentiation of hematopoietic cells.

Subcellular Localization: [Isoform 1]: Cell membrane; Single-pass type I membrane protein.; [Isoform 2]: Secreted.

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer-composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.