

Recombinant Mouse IL20 Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP52106

Size: 10ug, 100ug

Protein Name: Interleukin-20 (IL-20) (Cytokine Zcyto10)

Synonym: IL20, Zcyto10

UniProt ID: Q9JKV9

UniProt Link: www.uniprot.org/uniprotkb/Q9JKV9/entry

Species Source: Mouse

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 176 AA

Molecular Weight: 20 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MKGFGLAFLGLFSAVGFLLWTPLTGLKTLHLGSCVITANLQAIQKEFSEIRDSVQAEDTNIDIRILRTTES
LKDIKSLDRCCFLRHLVRFYLDLVFKVYQTPDHHTLRKISSLANSLFIKKDLSVCHSHMACHCGEEA
MEKYNQILSHFIELELQAAVVKALGELGILLRWMEEML

3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.

Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant Interleukin-20 (IL-20) (Cytokine Zcyto10) produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: Pro-inflammatory and angiogenic cytokine mainly secreted by monocytes and skin keratinocytes that plays crucial roles in immune responses, regulation of inflammatory responses, hemopoiesis, as well as epidermal cell and keratinocyte differentiation (PubMed:23793061). Enhances tissue remodeling and wound-healing activities and restores the homeostasis of epithelial layers during infection and inflammatory responses to maintain tissue integrity. Affects multiple actin-mediated functions in activated neutrophils leading to inhibition of phagocytosis, granule exocytosis, and migration. Exert its effects via the type I IL-20 receptor complex consisting of IL20RA and IL20RB. Alternatively, can mediate its activity through a second receptor complex called type II IL-20 receptor complex composed of IL22RA1 and IL20RB (By similarity). Acts as an arteriogenic and vascular remodeling factory by activating a range of signaling processes including phosphorylations of JAK2 and STAT5 as well as activation of the serine and threonine kinases AKT and ERK1/2 (PubMed:17878297). Alternatively, can activate STAT3 phosphorylation and transcriptional activity in a JAK2, ERK1/2 and p38 MAPK-dependent manner in keratinocytes (By similarity).

Subcellular Localization: Secreted.

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer-composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.