

Recombinant Mouse I10R2 Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP52058

Size: 10ug, 100ug

Protein Name: Interleukin-10 receptor subunit beta (IL-10 receptor subunit beta) (IL-10R subunit beta) (IL-10RB) (Cytokine receptor class-II member 4) (Cytokine receptor family 2 member 4) (CRF2-4) (Interleukin-10 receptor subunit 2) (IL-10R subunit 2) (IL-10R2) (CD antigen CDw210b)

Synonym: Il10rb, Crfb4

UniProt ID: Q61190

UniProt Link: www.uniprot.org/uniprotkb/Q61190/entry

Species Source: Mouse

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 349 AA

Molecular Weight: 40 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MAPCVAGWLGGFLLVPALGMIPPPEKVRMNSVNFKNILQWEVPAFPKTNLTFTAQYESYRSFQDHC
KRTASTQCDFSHLSKYGDYTVRVRAELADEHSEWVNVTFCPVEDTIIGPPEMQIESLAESLHLRFSA
PQIENEPETWTLKNIYDSWAYRVQYWKNGTNEKFQVVSYPYDSEVLRNLEPWTTYCIQVQGFLLDQN
RTGEWSEPICERTGNDEITPSWIVAILIVSVLVVFLFLLGCFVVLWLIYKKTKHTFRSGTSLPQHLKEFL
GHPHHSTFLLFSFPPPEEAEVFDKLSIIEESESESKQSPEDNCASEPPSDPGPRELESKDEAPSPPH
DDPKLLTSTSEV

3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.

Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant Interleukin-10 receptor subunit beta (IL-10 receptor subunit beta) (IL-10R subunit beta) (IL-10RB) (Cytokine receptor class-II member 4) (Cytokine receptor family 2 member 4) (CRF2-4) (Interleukin-10 receptor subunit 2) (IL-10R subunit 2) (IL-10R2) (CD antigen CDw210b) produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: Shared cell surface receptor required for the activation of five class 2 cytokines: IL10, IL22, IL26, IL28, and IFNL1. The IFNLR1/IL10RB dimer is a receptor for the cytokine ligands IFNL2 and IFNL3 and mediates their antiviral activity. The ligand/receptor complex stimulates the activation of the JAK/STAT signaling pathway leading to the expression of IFN-stimulated genes (ISG), which contribute to the antiviral state (By similarity).

Subcellular Localization: Membrane; Single-pass type I membrane protein.

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer-composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.