

Recombinant Human TGFR2 Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP50959

Size: 10ug, 100ug

Protein Name: TGF-beta receptor type-2 (TGFR-2) (EC 2.7.11.30) (TGF-beta type II receptor) (Transforming growth factor-beta receptor type II) (TGF-beta receptor type II) (TbetaR-II)

Synonym: TGFBR2

UniProt ID: P37173

UniProt Link: www.uniprot.org/uniprotkb/P37173/entry

Species Source: Human

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 567 AA

Molecular Weight: 65 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MGRGLLRGLWPLHIVLWTRIASTIPPHVQKSVNNDMIVTDNNGAVKFPQLCKFCDVRFSTCDNQKSC
MSNCSITSICEKPQEVCAVWRKNDENITLETVCHDPKLPYHDFILEDAAAPKCMKEKKKPGETFFM
CSCSSDECNDNIIFSEYNTSNPDLLLIVFQVTGISLLPPLGVAISVIIIIFYCYRVNRQQKLSSTWETGKT
RKLMEFSEHCAIILEDSDISSTCANNINHNTPELLDLVVGKGRFAEVYKAKLKQNTSEQFETVA
VKIFPYEEYASWKTEKDIFSDINLKHENILQFLTAEERKTEL GKQYWLITAFHAKGNLQEYLTRHVISW
EDLRKLGSSSLARGIAHLHSDHTPCGRPKMPIVHRDLKSSNILVKNDLTCCLCDFGLSLRLDPTLSVDD
LANSGQVGTARYMAPEVLESRMNLENVESFKQTDVYSMALVLWEMTSRCNAVGEVKDYEPFPGSK
VREHPCVESMKDNVLRDRGRPEIPSWLNLHQQIQMVCETLTECWDHDPEARLTAQCVAERFSELE
HLDRLSGRSCSEEKIPEDGSLNTTK

3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.

Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant TGF-beta receptor type-2 (TGFR-2) (EC 2.7.11.30) (TGF-beta type II receptor) (Transforming growth factor-beta receptor type II) (TGF-beta receptor type II) (TbetaR-II) produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: Transmembrane serine/threonine kinase forming with the TGF-beta type I serine/threonine kinase receptor, TGFBR1, the non-promiscuous receptor for the TGF-beta cytokines TGFB1, TGFB2 and TGFB3. Transduces the TGFB1, TGFB2 and TGFB3 signal from the cell surface to the cytoplasm and thus regulates a plethora of physiological and pathological processes including cell cycle arrest in epithelial and hematopoietic cells, control of mesenchymal cell proliferation and differentiation, wound healing, extracellular matrix production, immunosuppression and carcinogenesis. The formation of the receptor complex composed of 2 TGFBR1 and 2 TGFBR2 molecules symmetrically bound to the cytokine dimer results in the phosphorylation and activation of TGFBR1 by the constitutively active TGFBR2. Activated TGFBR1 phosphorylates SMAD2 which dissociates from the receptor and interacts with SMAD4. The SMAD2-SMAD4 complex is subsequently translocated to the nucleus where it modulates the transcription of the TGF-beta-regulated genes. This constitutes the canonical SMAD-dependent TGF-beta signaling cascade. Also involved in non-canonical, SMAD-independent TGF-beta signaling pathways.; [Isoform 1]: Has transforming growth factor beta-activated receptor activity.; [Isoform 2]: Has transforming growth factor beta-activated receptor activity.; [Isoform 3]: Binds TGFB1, TGFB2 and TGFB3 in the picomolar affinity range without the participation of additional receptors. Blocks activation of SMAD2 and SMAD3 by TGFB1.

Subcellular Localization: Cell membrane ; Single-pass type I membrane protein . Membrane raft .; [Isoform 3]: Secreted .

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer-composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.