

Recombinant Rat E2AK2 Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP52492

Size: 10ug, 100ug

Protein Name: Interferon-induced, double-stranded RNA-activated protein kinase (EC 2.7.11.1) (Eukaryotic translation initiation factor 2-alpha kinase 2) (eIF-2A protein kinase 2) (Interferon-inducible RNA-dependent protein kinase) (Protein kinase RNA-activated) (PKR) (Protein kinase R) (Tyrosine-protein kinase EIF2AK2) (EC 2.7.10.2)

Synonym: Eif2ak2, Prkr

UniProt ID: Q63184

UniProt Link: www.uniprot.org/uniprotkb/Q63184/entry

Species Source: Rat

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 513 AA

Molecular Weight: 58 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MASDTPGFYVDKLNKYSQIHKVKIYKEISVTGPPHRRFTFQVIIIEEREFPEGEGRSKQEAKNNAAKL
AVEILDNENKVDSDTDASEQGLIEGNYIGLVNSFAQKENLPVNFELCDPDSQLPHRFICKCKIGQTTY
GTGFGANKKEAKQLAAKNAYQKLSEKSPSKTGFTVSLSSDFSSSSSITSNSASQSASGRDFEDIFMN
GLREKRKSGVKVPSDDVLRNKYTLDDRFSKDFEDIEEIGSGGFGQVFKAKHRIDGKTYAIKRITYNTK
KAKREVQALAE LN HANIVQYRVCWEGEDYDYPENSTNGDTSRYKTRCLFIQMEFCDKGTLLQQWL
EKRNRSQEDKALVLELFEQIVTGVDYIHSKGLIHRDLKPGNIFLVDEKHIKIGDFGLATALENDGNPRT
KYTGTPQYMSPEQKSSSLVEYGKEVDIFALGLILAELLHICKTDSEKIEFFQLLRNGIFSDDIFDNKEKSL
LQKLLSSKPRERPNTSEILKTLAEWKNISEKKRNTC

3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.



Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant Interferon-induced, double-stranded RNA-activated protein kinase (EC 2.7.11.1) (Eukaryotic translation initiation factor 2-alpha kinase 2) (eIF-2A protein kinase 2) (Interferon-inducible RNA-dependent protein kinase) (Protein kinase RNA-activated) (PKR) (Protein kinase R) (Tyrosine-protein kinase EIF2AK2) (EC 2.7.10.2) produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: IFN-induced dsRNA-dependent serine/threonine-protein kinase that phosphorylates the alpha subunit of eukaryotic translation initiation factor 2 (EIF2S1/eIF-2-alpha) and plays a key role in the innate immune response to viral infection (By similarity). Inhibits viral replication via the integrated stress response (ISR): EIF2S1/eIF-2-alpha phosphorylation in response to viral infection converts EIF2S1/eIF-2-alpha in a global protein synthesis inhibitor, resulting to a shutdown of cellular and viral protein synthesis, while concomitantly initiating the preferential translation of ISR-specific mRNAs, such as the transcriptional activator ATF4 (By similarity). Exerts its antiviral activity on a wide range of DNA and RNA viruses (By similarity). Also involved in the regulation of signal transduction, apoptosis, cell proliferation and differentiation: phosphorylates other substrates including p53/TP53, PPP2R5A, DHX9, ILF3 and IRS1 (By similarity). In addition to serine/threonine-protein kinase activity, also has tyrosine-protein kinase activity and phosphorylates CDK1 at 'Tyr-4' upon DNA damage, facilitating its ubiquitination and proteasomal degradation (By similarity). Either as an adapter protein and/or via its kinase activity, can regulate various signaling pathways (p38 MAP kinase, NF-kappa-B and insulin signaling pathways) and transcription factors (JUN, STAT1, STAT3, IRF1, ATF3) involved in the expression of genes encoding pro-inflammatory cytokines and IFNs (By similarity). Activates the NF-kappa-B pathway via interaction with IKBKB and TRAF family of proteins and activates the p38 MAP kinase pathway via interaction with MAP2K6 (By similarity). Can act as both a positive and negative regulator of the insulin signaling pathway (ISP) (By similarity). Negatively regulates ISP by inducing the inhibitory phosphorylation of insulin receptor substrate 1 (IRS1) at 'Ser-312' and positively regulates ISP via phosphorylation of PPP2R5A which activates FOXO1, which in turn up-regulates the expression of insulin receptor substrate 2 (IRS2) (By similarity). Can regulate NLRP3 inflammasome assembly and the activation of NLRP3, NLRP1, AIM2 and NLRC4 inflammasomes (By similarity). Plays a role in the regulation of the cytoskeleton by binding to gelsolin (GSN), sequestering the protein in an inactive conformation away from actin (By similarity).

Subcellular Localization: Cytoplasm . Nucleus . Cytoplasm, perinuclear region .

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer-composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.