

Recombinant Mouse I13R2 Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP51199

Size: 10ug, 100ug

Protein Name: Interleukin-13 receptor subunit alpha-2 (IL-13 receptor subunit alpha-2) (IL-13R subunit alpha-2) (IL-13R-alpha-2) (IL-13RA2) (CD antigen CD213a2)

Synonym: Il13ra2

UniProt ID: O88786

UniProt Link: www.uniprot.org/uniprotkb/O88786/entry

Species Source: Mouse

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 383 AA

Molecular Weight: 44 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MAFVHIRCLCFILLCTITGYSLEIKVNPPQDFEILDPGLLGYLEYLQWKPPVIEKFKGCTLEYELKYRNV
DSDSWKTIITRNLIYKDGFDLNKGIEGKIRTHLSEHCTNGSEVQSPWIEASYGISDEGSLETKIQDMKCI
YYNWQYLVCWSWKPGKTVYSDTNYTMFFWYEGLDHALQCADYLQHDEKNVGCKLSNLDSSDYKDFF
ICVNGSSKLEPIRSSYTVFQLQNIVKPLPPEFLHISVENSIDIRMKWSTPGGPIPPRCYTYEIVIREDDIS
WESATDKNDMCLKRRANESEDL CFFVRCKVNIYCADDGIWSEWSEEECWEGYTGPDSKIIFIVPVCL
FFIFLLLLLCLIVEKEEPEPTLSLHVDLNKEVCAYEDTLC

3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.

Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant Interleukin-13 receptor subunit alpha-2 (IL-13 receptor subunit alpha-2) (IL-13R subunit alpha-2) (IL-13R-alpha-2) (IL-13RA2) (CD antigen CD213a2) produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: Cell surface receptor that plays a role in the regulation of IL-13-mediated responses (PubMed:12642602). Functions as a decoy receptor that inhibits IL-13- and IL-4-mediated signal transduction via the JAK-STAT pathway and thereby modulates immune responses and inflammation. Serves as a functional signaling receptor for IL-13 in an alternative pathway involving AP-1 ultimately leading to the production of TGFB1.

Subcellular Localization: [Isoform 1]: Cell membrane ; Single-pass type I membrane protein .; [Isoform 2]: Secreted.

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer-composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.