

Recombinant Mouse I15RA Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP51624

Size: 10ug, 100ug

Protein Name: Interleukin-15 receptor subunit alpha (IL-15 receptor subunit alpha) (IL-15R-alpha) (IL-15RA) (CD antigen CD215) [Cleaved into: Soluble interleukin-15 receptor subunit alpha (sIL-15 receptor subunit alpha) (sIL-15R-alpha) (sIL-15RA)]

Synonym: Il15ra

UniProt ID: Q60819

UniProt Link: www.uniprot.org/uniprotkb/Q60819/entry

Species Source: Mouse

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 263 AA

Molecular Weight: 28 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MASPQLRGYGVQAIPVLLLLLLLLLLLLPLRVTPGTTCPPPVSIEHADIRVKNYSVNSRERYVCNSGFKR
KAGTSTLIECVINKNTNVAHWTTPSLKCIRDPSLAHYSPVPTVTPKVTSQPESPSPSAKEPEAFSPK
SDTAMTTETAIMPGRSLTPSQTTSAGTTGTGSHKSSRAPSLAATMTLEPTASTSLRITEISPHSSKMT
KVAISTSVLLVGAGVMAFLAWYIKSRQPSQPCRVEVETMETVPMTVRASSKEDEDTGA

3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.

Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant Interleukin-15 receptor subunit alpha (IL-15 receptor subunit alpha) (IL-15R-alpha) (IL-15RA) (CD antigen CD215) [Cleaved into: Soluble interleukin-15 receptor subunit alpha (sIL-15 receptor subunit alpha) (sIL-15R-alpha) (sIL-15RA)] produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: High-affinity receptor for interleukin-15 (PubMed:17947230). Can signal both in cis and trans where IL15R from one subset of cells presents IL15 to neighboring IL2RG-expressing cells (PubMed:17947230). In neutrophils, binds and activates kinase SYK in response to IL15 stimulation (By similarity). In neutrophils, required for IL15-induced phagocytosis in a SYK-dependent manner (By similarity).

Subcellular Localization: Membrane ; Single-pass type I membrane protein . Nucleus membrane ; Single-pass type I membrane protein . Cell surface .; [Soluble interleukin-15 receptor subunit alpha]: Secreted, extracellular space .

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer-composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.