

Recombinant Mouse TNR8 Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP51626

Size: 10ug, 100ug

Protein Name: Tumor necrosis factor receptor superfamily member 8 (CD30L receptor)
(Lymphocyte activation antigen CD30) (CD antigen CD30)

Synonym: Tnfrsf8

UniProt ID: Q60846

UniProt Link: www.uniprot.org/uniprotkb/Q60846/entry

Species Source: Mouse

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 498 AA

Molecular Weight: 53 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MSALLTAAGLLFLGMLQAFPTDRPLKTTTCAGDLSHYPGEAARNCCYQCPSGLSPTQPCPRGPAHCR
KQCAPDYVYVNEDEGKCTACVTCLPGLVEKAPCSGNSPRICECQPGMHCCCTPAVNSCARCKLHCSGE
EYVVKSPGTAKKDTICELPSSSGSGPNCSNPGDRKTLTSHATPQAMPTLESPANDSARSLPMPRVTNL
VQEDATELVKVPESSSSKAREPSPDPGNAEKNMTLELPSPGTLPDISTSENSKEPASTASTLSLVVD
AWTSSRMQPTSPLSTGTPFLDPGPVLFVWAMVLLVGSGSFLLCYWKACRRRFQKFLDYLVT
FQPKMEQTDSCPTKLTQPQRSGSVTDPSSTGHKLSPVSPPPAVETCASVGATYLENLPLDDSPAG
NPFSPREPPEPRVSTEHTNNRIEKIYIMKADTVIVGSVKTEVPEGRAPAGSTESELEAELEVDHAPHY
PEQETEPPLGSCTEVMFSVEEGGKEDHGPTTVSEK

3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.

Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant Tumor necrosis factor receptor superfamily member 8 (CD30L receptor) (Lymphocyte activation antigen CD30) (CD antigen CD30) produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: Receptor for TNFSF8/CD30L (PubMed:8543792). May play a role in the regulation of cellular growth and transformation of activated lymphoblasts. Regulates gene expression through activation of NF-kappa-B (By similarity).

Subcellular Localization: Cell membrane ; Single-pass type I membrane protein .

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer-composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.