

Recombinant Human TPO Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP51008

Size: 10ug, 100ug

Protein Name: Thrombopoietin (C-mpl ligand) (ML) (Megakaryocyte colony-stimulating factor) (Megakaryocyte growth and development factor) (MGDF) (Myeloproliferative leukemia virus oncogene ligand)

Synonym: THPO, MGDF

UniProt ID: P40225

UniProt Link: www.uniprot.org/uniprotkb/P40225/entry

Species Source: Human

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 353 AA

Molecular Weight: 38 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MELTELLLVVMLLLTARLTLSPPACDLRVLSKLLRDSHVLHSRLSQCEVHPLPTPVLLPAVDFSL
GEWKTQMEETKAQDILGAVTLLLEGVMAARGQLGPTCLSSLLGQLSGQVRLLLGALQSLLGTQLPP
QGRRTAHKDPNAIFLSFQHLLRGKVRFLMLVGGSTLCVRRAPPTTAVPSRTSLVLTNLNLPNRTSGLL
ETNFTASARTTGSGLLKWQQGFRAKIPGLLNQTSRSLDQIPGYLNRIHELLNGTRGLFPGPSRRTL
APDISSGTSDTGSLPPNLQPGYSPSPHPPTGQYTLFPLPPTLPTPVVQLHPLLPDPSAPTPTPTSPL
LNTSYTHSQNLSQEG

3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.

Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant Thrombopoietin (C-mpl ligand) (ML) (Megakaryocyte colony-stimulating factor) (Megakaryocyte growth and development factor) (MGDF) (Myeloproliferative leukemia virus oncogene ligand) produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: Lineage-specific cytokine affecting the proliferation and maturation of megakaryocytes from their committed progenitor cells. It acts at a late stage of megakaryocyte development. It may be the major physiological regulator of circulating platelets.

Subcellular Localization: Secreted .

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer-composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.