

Recombinant Rat PDGFC Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP52571

Size: 10ug, 100ug

Protein Name: Platelet-derived growth factor C (PDGF-C) (Fallotin) (Spinal cord-derived growth factor) (rScdgc) (VEGF-E) [Cleaved into: Platelet-derived growth factor C, latent form (PDGFC latent form); Platelet-derived growth factor C, receptor-binding form (PDGFC receptor-binding form)]

Synonym: Pdgfc, Scdgc

UniProt ID: Q9EQX6

UniProt Link: www.uniprot.org/uniprotkb/Q9EQX6/entry

Species Source: Rat

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 345 AA

Molecular Weight: 39 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MLLLGLLLLTSALAGQRTGTRAESNLSSKLQLSSDKEQNGVQDPRHERVVTISGNGSIHSPKFPHTY
PRNTVLVWRLVAVDENVRIQLTFDERFGLEDPEDDLCKYDFVEVEEPSDGSVLGRWCGSGTVPGK
QTSKGNHIRIRFVSDEYFPSEPGFCIHYSIIMPQVTETTSPSVLPPSALSLLNNAVTAFSTVEELIRF
LEPDRWQIDLSLYKPTWPLLGAFLYGKKS KAVNLNLLKEEVKLYSCTPRNFSVSIREEELKRTDTIF
WPGCLLVKRCGGNCACCLHNCNECQCVPKVTKKYHEVLQLRPKIGVKGLHKS LTDVALEHHEEC
DCVCRGNTG

3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.

Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant Platelet-derived growth factor C (PDGF-C) (Fallotin) (Spinal cord-derived growth factor) (rScdrg) (VEGF-E) [Cleaved into: Platelet-derived growth factor C, latent form (PDGFC latent form); Platelet-derived growth factor C, receptor-binding form (PDGFC receptor-binding form)] produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: Growth factor that plays an essential role in the regulation of embryonic development, cell proliferation, cell migration, survival and chemotaxis. Potent mitogen and chemoattractant for cells of mesenchymal origin. Required for normal skeleton formation during embryonic development, especially for normal development of the craniofacial skeleton and for normal development of the palate. Required for normal skin morphogenesis during embryonic development. Plays an important role in wound healing, where it appears to be involved in three stages: inflammation, proliferation and remodeling. Plays an important role in angiogenesis and blood vessel development. Involved in fibrotic processes, in which transformation of interstitial fibroblasts into myofibroblasts plus collagen deposition occurs. The CUB domain has mitogenic activity in coronary artery smooth muscle cells, suggesting a role beyond the maintenance of the latency of the PDGF domain. In the nucleus, PDGFC seems to have additional function (By similarity).

Subcellular Localization: Cytoplasm, cytosol . Secreted . Nucleus . Cytoplasmic granule . Cell membrane . Note=Sumoylated form is predominant in the nucleus. Stored in alpha granules in platelets.

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer-composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.