

Recombinant Mouse EDAR Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP51975

Size: 10ug, 100ug

Protein Name: Tumor necrosis factor receptor superfamily member EDAR (Anhidrotic ectodysplasin receptor 1) (Downless) (Ectodermal dysplasia receptor) (Ectodysplasin-A receptor)

Synonym: Edar, DI

UniProt ID: Q9R187

UniProt Link: www.uniprot.org/uniprotkb/Q9R187/entry

Species Source: Mouse

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 448 AA

Molecular Weight: 48 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MAHVGDCCKWMSWLPVLVVSMLCSAKAEDSNCGENEYHNQTTGLCQQCPPCRPGEEPYMSCGYG
TKDDDYGCVPCPAEKFSKGGYQICRRHKDCEGFFRATVLTTPGDMENDAECGPCLPGYYMLENRPR
NIYGMVCYSCLLAPPNTKECVGATSGVSAHSSSTSGGSTLSFPQHAKELSGQGHLATALIIMSTIFI
MAIAIVLIIMFYIMKTKPSAPACCSSPPGKSAEAPANTHEEKKEAPDSVVTFPENGEFQKLTATPTKTP
KSENDASSENEQLLSRSVDSDEEPAPDKQGSPCLLSLVHLAREKSVTSNKSAGIQSRRKKILDVY
ANVCGVVEGLSPTELPFDCLKTSRMLSSTYNSEKAVVKTWRHLAESFGLKRDEIGGMTDGMQLFD
RISTAGYSIPELLTKLVQIERLDAVESLCADILEWAGVPPASPPPAAS

3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.

Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant Tumor necrosis factor receptor superfamily member EDAR (Anhidrotic ectodysplasin receptor 1) (Downless) (Ectodermal dysplasia receptor) (Ectodysplasin-A receptor) produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: Receptor for EDA isoform TAA, but not for EDA isoform TA-2 (By similarity). May mediate the activation of NF-kappa-B and JNK. May promote caspase-independent cell death.

Subcellular Localization: Membrane ; Single-pass type I membrane protein .

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer-composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.