

Recombinant Rat ATPO Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP52450

Size: 10ug, 100ug

Protein Name: ATP synthase peripheral stalk subunit OSCP, mitochondrial (ATP synthase subunit O) (Oligomycin sensitivity conferral protein) (OSCP) (Sperm flagella protein 4)

Synonym: Atp5po, Atp5o

UniProt ID: Q06647

UniProt Link: www.uniprot.org/uniprotkb/Q06647/entry

Species Source: Rat

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 213 AA

Molecular Weight: 23 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MAAPATSVLSRQVRSFSTSVVRPFSLVRPPVQVYGIEGRYATALYSAASKQKRLDQVEKELLRVG
QLLKDPKVSLAVLNPIYIKRSIKVKSLKDITTEKFSPILTANLMNLLAENGR LGNTQGVISAFSTIMSVHR
GEVPCTVTTAFPLDEAVLSELKTVLNSFLSKGQILNLEVKTDP SIMGGMIVRIGEKYVDMSAKSKIQKL
SKAMRDLL

3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.

Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant ATP synthase peripheral stalk subunit OSCP, mitochondrial (ATP synthase subunit O) (Oligomycin sensitivity conferral protein) (OSCP) (Sperm flagella protein 4) produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: Subunit OSCP, of the mitochondrial membrane ATP synthase complex (F(1)F(0) ATP synthase or Complex V) that produces ATP from ADP in the presence of a proton gradient across the membrane which is generated by electron transport complexes of the respiratory chain. ATP synthase complex consist of a soluble F(1) head domain - the catalytic core - and a membrane F(1) domain - the membrane proton channel. These two domains are linked by a central stalk rotating inside the F(1) region and a stationary peripheral stalk. During catalysis, ATP synthesis in the catalytic domain of F(1) is coupled via a rotary mechanism of the central stalk subunits to proton translocation (By similarity). In vivo, can only synthesize ATP although its ATP hydrolase activity can be activated artificially in vitro (By similarity). Part of the complex F(0) domain (By similarity). Part of the complex F(0) domain and the peripheric stalk, which acts as a stator to hold the catalytic alpha(3)beta(3) subcomplex and subunit a/ATP6 static relative to the rotary elements (By similarity).

Subcellular Localization: Mitochondrion . Mitochondrion inner membrane .

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer- composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.