

Recombinant Rat KPCD Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP52257

Size: 10ug, 100ug

Protein Name: Protein kinase C delta type (EC 2.7.11.13) (nPKC-delta) [Cleaved into: Protein kinase C delta type regulatory subunit; Protein kinase C delta type catalytic subunit (Sphingosine-dependent protein kinase-1) (SDK1)]

Synonym: Prkcd, Pkcd

UniProt ID: P09215

UniProt Link: www.uniprot.org/uniprotkb/P09215/entry

Species Source: Rat

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 673 AA

Molecular Weight: 78 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MAPFLRISFNSYELGSLQAEDDASQPFCAVKMKEALTDRGKTLVQKKPTMYPEWKSTFDAHIYEG
RVIQIVLMRAAEDPMSEVTGVSVLAERCKKNNGKAEFWLDLQPQAKVLMCVQYFLEDGDCKQSM
RSEEEAMFPTMNRRAIKQAKIHAIKNEFIATFFGQPTFCVCKEFVWGLNKQGYKCRQCNAAIHK
KCIDKIIGRCTGTATNSRDTIFQKERFNIDMPHRFKVYNYMSPTFCDHCGTLLWGLVKQGLKCEDCG
MNVHHKCREKVANLCGINQKLLAEALNQVTQKASRKPETPETVGIYQGFEKKTAVSGNDIPDNNGTY
GKIWEGSNRCLENFTFQKVLGKGSFGKVLAEELGKERYFAIKYLKKDVLIDDDVECTMVEKRVL
ALAWENPFLTHLICTFQTKDHLFFVMEFLNGGDLMFHIQDKGRFELYRATFYAAEIIICGLQFLHGKGI
YRDLKLDNVMLDKDGHIAIDFGMCKENIFGENRASTFCGTPDYIAPEILQGLKYSFSVDWWSFGVLL
YEMLIQSPFHGDDDELFEISIRVDTPHYPRWITKESKDIMEKLFERDPAKRLGVTGNIRLHPFFKTIN
WNLLEKRKVEPPFKPKVKSPSDYSNFDPEFLNEKPQLSFSDKNLIDSMQDTAFKGFVFNPKYEQFL
E

3. Buffer Composition



Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.

Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant Protein kinase C delta type (EC 2.7.11.13) (nPKC-delta) [Cleaved into: Protein kinase C delta type regulatory subunit; Protein kinase C delta type catalytic subunit (Sphingosine-dependent protein kinase-1) (SDK1)] produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: Calcium-independent, phospholipid- and diacylglycerol (DAG)-dependent serine/threonine-protein kinase that plays contrasting roles in cell death and cell survival by functioning as a pro-apoptotic protein during DNA damage-induced apoptosis, but acting as an anti-apoptotic protein during cytokine receptor-initiated cell death, is involved in tumor suppression, is required for oxygen radical production by NADPH oxidase and acts as a positive or negative regulator in platelet functional responses. Upon DNA damage, activates the promoter of the death-promoting transcription factor BCLAF1/Btf to trigger BCLAF1-mediated p53/TP53 gene transcription and apoptosis. In response to oxidative stress, interact with and activate CHUK/IKKA in the nucleus, causing the phosphorylation of p53/TP53. In the case of ER stress or DNA damage-induced apoptosis, can form a complex with the tyrosine-protein kinase ABL1 which trigger apoptosis independently of p53/TP53. In cytosol can trigger apoptosis by activating MAPK11 or MAPK14, inhibiting AKT1 and decreasing the level of X-linked inhibitor of apoptosis protein (XIAP), whereas in nucleus induces apoptosis via the activation of MAPK8 or MAPK9. Upon ionizing radiation treatment, is required for the activation of the apoptosis regulators BAX and BAK, which trigger the mitochondrial cell death pathway. Can phosphorylate MCL1 and target it for degradation which is sufficient to trigger for BAX activation and apoptosis. Is required for the control of cell cycle progression both at G1/S and G2/M phases. Mediates phorbol 12-myristate 13-acetate (PMA)-induced inhibition of cell cycle progression at G1/S phase by up-regulating the CDK inhibitor CDKN1A/p21 and inhibiting the cyclin CCNA2 promoter activity. In response to UV irradiation can phosphorylate CDK1, which is important for the G2/M DNA damage checkpoint activation. Can protect glioma cells from the apoptosis induced by TNFSF10/TRAIL, probably by inducing increased phosphorylation and subsequent activation of AKT1. Can also act as tumor suppressor upon mitogenic stimulation with PMA or TPA. In N-formyl-methionyl-leucyl-phenylalanine (fMLP)-treated cells, is required for NCF1 (p47-phox) phosphorylation and activation of NADPH oxidase activity, and regulates TNF-elicited superoxide anion production in neutrophils, by direct phosphorylation and activation of NCF1 or indirectly through MAPK1/3 (ERK1/2) signaling pathways. Involved in antifungal immunity by mediating phosphorylation and activation of CARD9 downstream of C-type lectin receptors activation, promoting interaction between CARD9 and BCL10, followed by activation of NF-kappa-B and MAP kinase p38 pathways (By similarity). May also play a role in the regulation of NADPH oxidase activity in eosinophil after stimulation with IL5, leukotriene B4 or PMA. In collagen-induced platelet aggregation, acts a negative regulator of filopodia formation and actin polymerization by interacting with and negatively regulating VASP phosphorylation. Downstream of PAR1, PAR4 and CD36/GP4 receptors, regulates differentially platelet dense granule secretion; acts as a positive regulator in PAR-mediated granule secretion, whereas it negatively regulates CD36/GP4-mediated granule release. Phosphorylates MUC1 in the C-terminal and regulates the interaction between

MUC1 and beta-catenin (By similarity). The catalytic subunit phosphorylates 14-3-3 proteins (YWHAB, YWHAZ and YWHAH) in a sphingosine-dependent fashion. Phosphorylates ELAVL1 in response to angiotensin-2 treatment (By similarity). Phosphorylates mitochondrial phospholipid scramblase 3 (PLSCR3), resulting in increased cardiolipin expression on the mitochondrial outer membrane which facilitates apoptosis (By similarity). Phosphorylates SMPD1 which induces SMPD1 secretion (By similarity). Acts as a negative regulator of smoothened signaling by mediating phosphorylation of GLI1, preventing its localization to the nucleus (By similarity). Acts as a regulator of dopamine signaling by mediating constitutive phosphorylation of dopamine receptor DRD1, inhibiting G protein-coupled receptor signaling (By similarity).; Truncated isoform 2 is inactive.

Subcellular Localization: Cytoplasm . Nucleus . Cytoplasm, perinuclear region . Cell membrane ; Peripheral membrane protein . Mitochondrion . Endomembrane system . Note=Translocates to the mitochondria upon apoptotic stimulation. Upon activation, translocates to the plasma membrane followed by partial location to the endolysosomes.

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer-composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.