

Recombinant Mouse MIF Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP51388

Size: 10ug, 100ug

Protein Name: Macrophage migration inhibitory factor (MIF) (EC 5.3.2.1) (Delayed early response protein 6) (DER6) (Glycosylation-inhibiting factor) (GIF) (L-dopachrome isomerase) (L-dopachrome tautomerase) (EC 5.3.3.12) (Phenylpyruvate tautomerase)

Synonym: Mif

UniProt ID: P34884

UniProt Link: www.uniprot.org/uniprotkb/P34884/entry

Species Source: Mouse

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 115 AA

Molecular Weight: 13 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MPMFIVNTNVPRASVPEGFLSELTQQLAQATGKPAQYIAVHVVPDQLMTFSGTNDPCALCSLHSIGKI
GGAQNRNYSKLLCGLLSDRLHISPDRVYINYYDMNAANVGWNGSTFA

3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.

Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant Macrophage migration inhibitory factor (MIF) (EC 5.3.2.1) (Delayed early response protein 6) (DER6) (Glycosylation-inhibiting factor) (GIF) (L-dopachrome isomerase) (L-dopachrome tautomerase) (EC 5.3.3.12) (Phenylpyruvate tautomerase) produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: Pro-inflammatory cytokine involved in the innate immune response to bacterial pathogens (By similarity). The expression of MIF at sites of inflammation suggests a role as mediator in regulating the function of macrophages in host defense (By similarity). Counteracts the anti-inflammatory activity of glucocorticoids (By similarity). Has phenylpyruvate tautomerase and dopachrome tautomerase activity (in vitro), but the physiological substrate is not known (PubMed:10933783, PubMed:16780921, PubMed:19188446). It is not clear whether the tautomerase activity has any physiological relevance, and whether it is important for cytokine activity (PubMed:10933783, PubMed:16780921, PubMed:19188446).

Subcellular Localization: Secreted . Cytoplasm . Note=Does not have a cleavable signal sequence and is secreted via a specialized, non-classical pathway. Secreted by macrophages upon stimulation by bacterial lipopolysaccharide (LPS), or by M.tuberculosis antigens (By similarity).

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer- composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.