

Recombinant Mouse BMP5 Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP51445

Size: 10ug, 100ug

Protein Name: Bone morphogenetic protein 5 (BMP-5)

Synonym: Bmp5, Bmp-5

UniProt ID: P49003

UniProt Link: www.uniprot.org/uniprotkb/P49003/entry

Species Source: Mouse

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 452 AA

Molecular Weight: 52 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MHWTVFLLRGIVGFLWSGWVQVGYAKGGLGDNHVHSSFIYRRLRNHERREIQREILSILGLPHRPRP
FSPGKQASSAPLFMLDLYNAMASEDNPEEYLVRVSLAGEAKETRKGYSPNGYAHRLHLPRTPL
TTQSPPLASLHDTNFLNDADMVMSFVNLVERDKDFSHQRRHYKEFRFDLTQIPHGEAVTAAEFRIYK
DKGNHRFENETIKISIIYQIIKEYTNRDADFLLDTRKTQALDVGWLVFDITVTSNHWVINPQNNLGLQL
CAETGDGRSINVKSAGLVGRHGPPQSKQPFMVAFFKASEVLLRSVRAASKRKNQNRNKSNSHQDPS
RMPSAGDYNTSEQKQACKKHELYVSFRDLGWQDWIIAPEGYAAFYCDGECFPLNAHNMNATNHAIV
QTLVHLMFPDHVPKPCCAPTKLNAISVLYFDDSSNVILKKYRNMVVRSCGCH

3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.

Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant Bone morphogenetic protein 5 (BMP-5) produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: Growth factor of the TGF-beta superfamily that plays essential roles in many developmental processes, including cartilage and bone formation or neurogenesis (PubMed:29321139, PubMed:7958439). Initiates the canonical BMP signaling cascade by associating with type I receptor BMPR1A and type II receptor BMPR2. In turn, BMPR1A propagates signal by phosphorylating SMAD1/5/8 that travel to the nucleus and act as activators and repressors of transcription of target genes. Can also signal through non-canonical pathway such as MAPK p38 signaling cascade to promote chondrogenic differentiation (By similarity). Promotes the expression of HAMP, this is repressed by its interaction with ERFE (PubMed:30097509).

Subcellular Localization: Secreted .

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer-composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.