

Recombinant Mouse BAD Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP51640

Size: 10ug, 100ug

Protein Name: Bcl2-associated agonist of cell death (BAD) (Bcl-2-binding component 6) (Bcl-xL/Bcl-2-associated death promoter) (Bcl2 antagonist of cell death)

Synonym: Bad, Bbc6

UniProt ID: Q61337

UniProt Link: www.uniprot.org/uniprotkb/Q61337/entry

Species Source: Mouse

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 204 AA

Molecular Weight: 22 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MGTPKQPSLAPAHALGLRKSDPGIRSLGSDAGGRRWRPAAQSMFQIPEFEPSEQEDASATDRGLG
PSLTEDQPGPYLAPGLLSNIHQQGRAATNSHHGGAGAMETRSHSSYPAGTEEDEGMEEELSPF
RGRSRSAPPNLWAAQRYGRELRRMSDEFEGSFKGLPRPKSAGTATQMRQSAGWTRIIQSWWDRN
LGKGGSTPSQ

3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.

Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant Bcl2-associated agonist of cell death (BAD) (Bcl-2-binding component 6) (Bcl-xL/Bcl-2-associated death promoter) (Bcl2 antagonist of cell death) produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: Promotes cell death. Successfully competes for the binding to Bcl-X(L), Bcl-2 and Bcl-W, thereby affecting the level of heterodimerization of these proteins with BAX. Can reverse the death repressor activity of Bcl-X(L), but not that of Bcl-2. Appears to act as a link between growth factor receptor signaling and the apoptotic pathways.

Subcellular Localization: Mitochondrion outer membrane. Cytoplasm . Note=Colocalizes with HIF3A isoform 2 in the cytoplasm (PubMed:21546903). Upon phosphorylation, locates to the cytoplasm.

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer-composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.