

Recombinant Rat HG2A Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP52267

Size: 10ug, 100ug

Protein Name: H-2 class II histocompatibility antigen gamma chain (Ia antigen-associated invariant chain) (Ii) (MHC class II-associated invariant chain) (CD antigen CD74) [Cleaved into: Class-II-associated invariant chain peptide (CLIP)]

Synonym: Cd74

UniProt ID: P10247

UniProt Link: www.uniprot.org/uniprotkb/P10247/entry

Species Source: Rat

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 280 AA

Molecular Weight: 32 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MDDQRDLISNHEQLPILGQRRAPESNCNRGBLYTSVSVLVALLAGQATTAYFLYQQQGRLDKLTV
TSQNLQLENLRMKLPKSAKPVSPMRMATPLLMRPLSMDNMLQAPVKNVTKYGNMTQDHVMHLLTK
SGPVNYPQLKGSFPENLKHKLNSMNGLDWKVFESWMKQWLLFEMSKNSLEEKQPTQTTPKVLTKC
QEEVSHIPDVHPGAFRPKCDENGNYMPLQCHGSTGYCWCVPNGTEVPHTKSRGRHNCSEPLDM
EDPSSGLGVTKQDMGQMFL

3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.

Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant H-2 class II histocompatibility antigen gamma chain (Ia antigen-associated invariant chain) (Ii) (MHC class II-associated invariant chain) (CD antigen CD74) [Cleaved into: Class-II-associated invariant chain peptide (CLIP)] produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: Plays a critical role in MHC class II antigen processing by stabilizing peptide-free class II alpha/beta heterodimers in a complex soon after their synthesis and directing transport of the complex from the endoplasmic reticulum to compartments where peptide loading of class II takes place. Enhance also the stimulation of T-cell responses through interaction with CD44.; [Class-II-associated invariant chain peptide]: Binds to the peptide-binding site of MHC class II alpha/beta heterodimers forming an alpha-beta-CLIP complex, thereby preventing the loading of antigenic peptides to the MHC class II complex until its release by HLA-DM in the endosome.; [Isoform Long]: Stabilizes the conformation of mature CTSL by binding to its active site and serving as a chaperone to help maintain a pool of mature enzyme in endocytic compartments and extracellular space of antigen-presenting cells (APCs).

Subcellular Localization: [Isoform Long]: Late endosome . Lysosome .; Cell membrane ; Single-pass type II membrane protein . Endoplasmic reticulum membrane . Golgi apparatus, trans-Golgi network . Endosome . Lysosome . Secreted . Note=Transits through a number of intracellular compartments in the endocytic pathway. It can either undergo proteolysis or reach the cell membrane.

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer-composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.