

Recombinant Mouse JUN Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP51230

Size: 10ug, 100ug

Protein Name: Transcription factor Jun (AH119) (Activator protein 1) (AP1) (Proto-oncogene c-Jun) (Transcription factor AP-1 subunit Jun) (V-jun avian sarcoma virus 17 oncogene homolog) (Jun A)

Synonym: Jun

UniProt ID: P05627

UniProt Link: www.uniprot.org/uniprotkb/P05627/entry

Species Source: Mouse

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 334 AA

Molecular Weight: 36 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MTAKMETTFYDDALNASFLQSESGAYGYSNPKILKQSMTLNLADPVGSLKPHLRAKNSDLLTSPDVG
LLKLASPELERLIIQSSNGHITTTPTPTQFLCPKNVTDEQEGFAEGFVRALAEHSQNTLPSVTSAAQP
VSGAGMVAPAVASVAGAGGGGGYSASLHSEPPVYANLSNPNPGALSSGGGAPSYGAAGLAFPSQ
PQQQQQPPQPPHLPQQIPVQHPRQLALKEEPQTVPEMPGETPPLSPIDMESQERIKAEKRMNRN
RIAASKCRKRLERLARLEEKVKTLKAQNSELASTANMLREQVAQLKQKVMNHVNSGCQLMLTQQL
QTF

3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.

Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant Transcription factor Jun (AH119) (Activator protein 1) (AP1) (Proto-oncogene c-Jun) (Transcription factor AP-1 subunit Jun) (V-jun avian sarcoma virus 17 oncogene homolog) (Jun A) produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: Transcription factor that recognizes and binds to the AP-1 consensus motif 5'-TGA[GC]TCA-3' (PubMed:14707112). Heterodimerizes with proteins of the FOS family to form an AP-1 transcription factor complex, thereby enhancing its DNA binding activity to the AP-1 consensus sequence 5'-TGA[GC]TCA-3' and enhancing its transcriptional activity (PubMed:2498083). Together with FOSB, plays a role in activation-induced cell death of T cells by binding to the AP-1 promoter site of FASLG/CD95L, and inducing its transcription in response to activation of the TCR/CD3 signaling pathway (By similarity). Promotes activity of NR5A1 when phosphorylated by HIPK3 leading to increased steroidogenic gene expression upon cAMP signaling pathway stimulation (PubMed:17210646). Involved in activated KRAS-mediated transcriptional activation of USP28 (By similarity). Binds to the USP28 promoter (By similarity).

Subcellular Localization: Nucleus .

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer-composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.