

Recombinant Human PDGFB Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP50773

Size: 10ug, 100ug

Protein Name: Platelet-derived growth factor subunit B (PDGF subunit B) (PDGF-2)
(Platelet-derived growth factor B chain) (Platelet-derived growth factor beta polypeptide)
(Proto-oncogene c-Sis) (Becaplermin)

Synonym: PDGFB, PDGF2, SIS

UniProt ID: P01127

UniProt Link: www.uniprot.org/uniprotkb/P01127/entry

Species Source: Human

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 241 AA

Molecular Weight: 27 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MNRCWALFLSLCCYLRLVSAEGDPIPEELYEMLSDHSIRSFDDLQRL LHGDPGEEDGAELDLNMTRS
HSGGELESLARGRRSLGSLTIAEPAMIAECKTRTEVFEISRRLIDRTNANFLVWPPCVEVQRCSGCC
NNRNQCRPTQVQLRPVQVRKIEIVRKKPIFKKATVTLEDHLACKCETVAAARPVTRSPGGSQEQRA
KTPQTRVTIRTVRVRPPKGKHKRKFHDKTALKETLGA

3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.

Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant Platelet-derived growth factor subunit B (PDGF subunit B) (PDGF-2) (Platelet-derived growth factor B chain) (Platelet-derived growth factor beta polypeptide) (Proto-oncogene c-Sis) (Becaplermin) produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: Growth factor that plays an essential role in the regulation of embryonic development, cell proliferation, cell migration, survival and chemotaxis. Potent mitogen for cells of mesenchymal origin (PubMed:26599395). Required for normal proliferation and recruitment of pericytes and vascular smooth muscle cells in the central nervous system, skin, lung, heart and placenta. Required for normal blood vessel development, and for normal development of kidney glomeruli. Plays an important role in wound healing. Signaling is modulated by the formation of heterodimers with PDGFA (By similarity).

Subcellular Localization: Secreted. Note=Released by platelets upon wounding.

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer-composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.