

Recombinant Rat CNOT9 Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP52474

Size: 10ug, 100ug

Protein Name: CCR4-NOT transcription complex subunit 9 (Cell differentiation protein RQCD1 homolog) (Rcd-1)

Synonym: Cnot9, Rcd1, Rqcd1

UniProt ID: Q5PQL2

UniProt Link: www.uniprot.org/uniprotkb/Q5PQL2/entry

Species Source: Rat

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 299 AA

Molecular Weight: 34 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MHSLATAAPVPTALAQVDREKIYQWINELSSPETRENALLELSKKRESVPDLAPMLWHSFGTIAALLQ
EIVNIYPSINPPTLTAHQSNRVCNALALLQCVASHPETRSAFLAAHIPLFLYPFLHTVSKTRPFYLRRLT
SLGVIGALVKTDQEVEINFLTTEIIPLCRIMESGSELSKTVATFILQKILLDDTGLAYICQTYERFSHVA
MILGKMVLQLSKEPSARLLKHVVRCYLRLSDNPRAREALRQCLPDQLKDDTTFAQVLKDDTTTKRWLA
QLVKNLQEGQVTDPRGIPLPPQ

3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.

Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant CCR4-NOT transcription complex subunit 9 (Cell differentiation protein RQCD1 homolog) (Rcd-1) produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: Component of the CCR4-NOT complex which is one of the major cellular mRNA deadenylases and is linked to various cellular processes including bulk mRNA degradation, miRNA-mediated repression, translational repression during translational initiation and general transcription regulation. Additional complex functions may be a consequence of its influence on mRNA expression. Involved in down-regulation of MYB- and JUN-dependent transcription. Enhances ligand-dependent transcriptional activity of nuclear hormone receptors. May play a role in cell differentiation (By similarity).

Subcellular Localization: Nucleus . Cytoplasm, P-body . Note=NANOS2 promotes its localization to P-body.

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer-composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.