

Recombinant Mouse RARA Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP51265

Size: 10ug, 100ug

Protein Name: Retinoic acid receptor alpha (RAR-alpha) (Nuclear receptor subfamily 1 group B member 1)

Synonym: Rara, Nr1b1

UniProt ID: P11416

UniProt Link: www.uniprot.org/uniprotkb/P11416/entry

Species Source: Mouse

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 462 AA

Molecular Weight: 51 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MASNSSSCPTPGGGHLNGYPVPPYAFFFFPMLGGLSPPGALTSLQHQLPVSGYSTPSPATIENTQSS
SSEEIVPSPSPPLPRIYKPCFVCQDKSSGYHYGVSACEGCKGFFRRSIQKNMYYTCHRDKNCIIN
KVTRNRCQYCRLQKCFDVGMSKESVRNDRNKKKKEAPKPECSESYTLTPEVGELIEKVRKAHQETF
PALCQLGKYTTNNSSEQRVSLDIDLWDKFSELSTKCIKTVEFAKQLPGFTTLTIADQITLLKAACLDILI
LRICTRYTPEQDTMTFSDGLTLNRTQMHNAGFGPLTDLVFAFANQLLPLEMDDAETGLLSAICLICGD
RQDLEQPDKVDMLQEPLLEALKVYVRKRRPSRPHMFPKMLMKITDLRSISAKGAERVITLKMEIPGS
MPPLIQEMLENSEGLDTLSGQSGGGTRDGGGLAPPPGSCSPSLSPSSHRSSPATQSP

3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.

Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant Retinoic acid receptor alpha (RAR-alpha) (Nuclear receptor subfamily 1 group B member 1) produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: Receptor for retinoic acid (PubMed:17205979). Retinoic acid receptors bind as heterodimers to their target response elements in response to their ligands, all-trans or 9-cis retinoic acid, and regulate gene expression in various biological processes (PubMed:17205979). The RXR/RAR heterodimers bind to the retinoic acid response elements (RARE) composed of tandem 5'-AGGTCA-3' sites known as DR1-DR5 (PubMed:17205979). In the absence of ligand, the RXR-RAR heterodimers associate with a multiprotein complex containing transcription corepressors that induce histone deacetylation, chromatin condensation and transcriptional suppression (By similarity). On ligand binding, the corepressors dissociate from the receptors and associate with the coactivators leading to transcriptional activation (PubMed:17205979, PubMed:19078967, PubMed:9230306). Formation of heterocomplex with histone deacetylases might lead to inhibition of RARE DNA element binding and to transcriptional repression (By similarity). Transcriptional activation and RARE DNA element binding might be supported by the transcription factor KLF2 (By similarity). RARA plays an essential role in the regulation of retinoic acid-induced germ cell development during spermatogenesis (PubMed:15901285). Has a role in the survival of early spermatocytes at the beginning prophase of meiosis (PubMed:15901285, PubMed:17905941). In Sertoli cells, may promote the survival and development of early meiotic prophase spermatocytes (PubMed:10660575, PubMed:17905941). In concert with RARG, required for skeletal growth, matrix homeostasis and growth plate function (PubMed:19389355). Together with RXRA, positively regulates microRNA-10a expression, thereby inhibiting the GATA6/VCAM1 signaling response to pulsatile shear stress in vascular endothelial cells (By similarity). In association with HDAC3, HDAC5 and HDAC7 corepressors, plays a role in the repression of microRNA-10a and thereby promotes the inflammatory response (By similarity).

Subcellular Localization: Nucleus . Cytoplasm . Note=Nuclear localization depends on ligand binding, phosphorylation and sumoylation (PubMed:10660575, PubMed:12079996). Translocation to the nucleus is dependent on activation of PKC and the downstream MAPK phosphorylation (PubMed:10660575, PubMed:12079996). Increased nuclear localization upon pulsatile shear stress (By similarity).

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer- composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.