

Recombinant Mouse IL25 Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP51810

Size: 10ug, 100ug

Protein Name: Interleukin-25 (IL-25) (Interleukin-17E) (IL-17E)

Synonym: Il25, Il17e

UniProt ID: Q8VHH8

UniProt Link: www.uniprot.org/uniprotkb/Q8VHH8/entry

Species Source: Mouse

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 169 AA

Molecular Weight: 19 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MYQAVAFLAMIVGTHTVSLRIQEGCSHLPSCCPSKEQEPPEEWLKWSSASVSPPEPLSHTTHHAESC
RASKDGPLNSRAISPWSYELDRDLNRVPQDLYHARCLCPHCVSLQTGSHMDPLGNSVPLYHNQTV
FYRRPCHGEEGTHRRYCLERRLYRVSLACVCVRPRVMA

3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.

Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant Interleukin-25 (IL-25) (Interleukin-17E) (IL-17E) produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: Cytokine produced by various cells such as eosinophils, T-helper type 2 (Th2) cells or epithelial cells that plays a role in internal safety of adaptive immune responses by regulating cytokine production (PubMed:11754819, PubMed:35615348). Promotes and augments T-helper type 2 responses locally or systemically (PubMed:35615348). Exerts its activity via its receptor composed of IL17RA and IL17RB for signal transduction (PubMed:18768888). In turn, stimulates the JAK2-STAT5A pathway and promotes the secretion of type-2 associated cytokines including IL4, IL9 and IL13. Also induces the release of IL8, and IL6 from eosinophils through the combined activation of MAPK and NF-kappa-B pathways. Inhibits the differentiation of T-helper (Th17) cells via the production of IL4, IL5 and IL13.

Subcellular Localization: Secreted .

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer- composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.