

Recombinant Mouse AIMP1 Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP51377

Size: 10ug, 100ug

Protein Name: Aminoacyl tRNA synthase complex-interacting multifunctional protein 1 (Multisynthase complex auxiliary component p43) [Cleaved into: Endothelial monocyte-activating polypeptide 2 (EMAP-2) (Endothelial monocyte-activating polypeptide II) (EMAP-II) (Small inducible cytokine subfamily E member 1)]

Synonym: Aimp1, Emap2, Scye1

UniProt ID: P31230

UniProt Link: www.uniprot.org/uniprotkb/P31230/entry

Species Source: Mouse

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 310 AA

Molecular Weight: 34 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MATNDAVLKRLEQKGAEADQII EYLKQQVALLKEKAILQATMREEKKLRVENAKLKKEIEELKQELILA
EIHNGVEQVRVRLSTPLQTNCTASESVVQSPSVATTASPATKEQIKAGEEKKVKEKTEKKGEKKEKQ
QSAAASTDSKPIDASRLDLRIGCIVTAKKHPDADSLYVEEVDVGEAAPRTVVSGLVNHVPLEQMQR
MVLNLCNLKPAKMRGVLSQAMVMCASSPEKVEILAPPNGSVPGDRITFDAFPGEKELNPKKKIWE
QIQPDLHTNAECVATYKGAPFEVKGKGVCRQAQTMANSIK

3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.

Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant Aminoacyl tRNA synthase complex-interacting multifunctional protein 1 (Multisynthase complex auxiliary component p43) [Cleaved into: Endothelial monocyte-activating polypeptide 2 (EMAP-2) (Endothelial monocyte-activating polypeptide II) (EMAP-II) (Small inducible cytokine subfamily E member 1)] produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: Non-catalytic component of the multisynthase complex (PubMed:12060739). Stimulates the catalytic activity of cytoplasmic arginyl-tRNA synthase (By similarity). Binds tRNA. Possesses inflammatory cytokine activity (PubMed:1400342, PubMed:7545917). Negatively regulates TGF-beta signaling through stabilization of SMURF2 by binding to SMURF2 and inhibiting its SMAD7-mediated degradation (PubMed:18448069). Involved in glucose homeostasis through induction of glucagon secretion at low glucose levels (PubMed:17001013). Promotes dermal fibroblast proliferation and wound repair (PubMed:15681823). Regulates KDELR1-mediated retention of HSP90B1/gp96 in the endoplasmic reticulum (PubMed:17525271). Plays a role in angiogenesis by inducing endothelial cell migration at low concentrations and endothelial cell apoptosis at high concentrations (By similarity). Induces maturation of dendritic cells and monocyte cell adhesion (PubMed:18292511). Modulates endothelial cell responses by degrading HIF-1A through interaction with PSMA7 (By similarity).

Subcellular Localization: Nucleus . Cytoplasm, cytosol . Secreted . Endoplasmic reticulum . Golgi apparatus . Note=Enriched in secretory vesicles of pancreatic alpha cells and secreted from the pancreas in response to low glucose levels (PubMed:17001013). Secreted in response to hypoxia (By similarity). Also secreted in response to both apoptotic and necrotic cell death.

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer-composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.