

Recombinant Mouse CD4 Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP51231

Size: 10ug, 100ug

Protein Name: T-cell surface glycoprotein CD4 (T-cell differentiation antigen L3T4) (T-cell surface antigen T4/Leu-3) (CD antigen CD4)

Synonym: Cd4

UniProt ID: P06332

UniProt Link: www.uniprot.org/uniprotkb/P06332/entry

Species Source: Mouse

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 457 AA

Molecular Weight: 51 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MCRAISLRRLLLLLLQLSQLLAVTQGKTLVLGKEGESAELPCESSQKKITVFTWKFSDQRKILGQH GK
GVLIRGGSPSQFDRFDSKKGAWEEKGSFPLIINKLMEDSQTICYELNRKEEVELWVFKVTFSPGTSL
LQGQSLTLTLDNSKVSNNPLTECKHKKGKVVS GSKVLSMSNLRVQDSDFWNCTVTLDQKKNWFGM
TL SVLG FQSTAITAYKSEGESA EFSFPLNFAEENGWGELMWKA EKDSFFQPWISFSIKNKEVSVQKS
TKDLKLQLKETLPLTLKIPQVSLQFAGSGNLT LTLDKGT LHQEVNLVVMKVAQLNNTLTCEVMGPTSP
KMRLTLKQENQEARVSEEQKV VQV VAPETGLWQCLLSEGD KVKMDSRIQVLSRGVNQTVFLACVL
GGSFGFLGFLGLCILCCVRCRHQQRQAARMSQIKRLLSEKKTCCPHRMQKSHNLI

3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.

Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant T-cell surface glycoprotein CD4 (T-cell differentiation antigen L3T4) (T-cell surface antigen T4/Leu-3) (CD antigen CD4) produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: Integral membrane glycoprotein that plays an essential role in the immune response and serves multiple functions in responses against both external and internal offenses. In T-cells, functions primarily as a coreceptor for MHC class II molecule:peptide complex. The antigens presented by class II peptides are derived from extracellular proteins while class I peptides are derived from cytosolic proteins. Interacts simultaneously with the T-cell receptor (TCR) and the MHC class II presented by antigen presenting cells (APCs). In turn, recruits the Src kinase LCK to the vicinity of the TCR-CD3 complex. LCK then initiates different intracellular signaling pathways by phosphorylating various substrates ultimately leading to lymphokine production, motility, adhesion and activation of T-helper cells. In other cells such as macrophages or NK cells, plays a role in differentiation/activation, cytokine expression and cell migration in a TCR/LCK-independent pathway. Participates in the development of T-helper cells in the thymus and triggers the differentiation of monocytes into functional mature macrophages.

Subcellular Localization: Cell membrane ; Single-pass type I membrane protein .

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer-composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.