

Recombinant Mouse GATA3 Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP51334

Size: 10ug, 100ug

Protein Name: Trans-acting T-cell-specific transcription factor GATA-3 (GATA-binding factor 3)

Synonym: Gata3, Gata-3

UniProt ID: P23772

UniProt Link: www.uniprot.org/uniprotkb/P23772/entry

Species Source: Mouse

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 443 AA

Molecular Weight: 48 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MEVTADQPRWVSHHHPAVLNGQHPDTHHPGLGHSYMEAQYPLTEEVDVLFNIDGQGNHVPSYYG
NSVRATVQRYPPTHHGSQVCRPPLLHGSLPWLDGGKALSSHHTASPWNLSPFSKTSIIHHGSPGPLS
VYPPASSSSLAAGHSSPHLFTFPPTPPKDVSPDPSLSTPGSAGSARQDEKECLKYQVQLPDSMKLE
TSHSRGSMTTLGASSSAHHPIITYPPYVPEYSSGLFPPSSLLGGSPTGFGCKSRPKARSSTEGRE
CVNCGATSTPLWRRDGTGHYLCNACGLYHKMNGQNRPLIKPKRRLSAARRAGTSCANCQTTTTTL
WRRNANGDPVCNACGLYYKLHNINRPLTMKKEGIQTRNRKMSSKSKKCKKVHDALEDFPKSSSFNP
AALSRHMSSLSHISPFSSHMLTTPTPMHPPSGLSFGPHHPSSMVTAMG

3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.

Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant Trans-acting T-cell-specific transcription factor GATA-3 (GATA-binding factor 3) produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: Transcriptional activator which binds to the enhancer of the T-cell receptor alpha and delta genes. Binds to the consensus sequence 5'-AGATAG-3'. Required for the T-helper 2 (Th2) differentiation process following immune and inflammatory responses. Positively regulates ASB2 expression (PubMed:31175139). Coordinates macrophage transcriptional activation and UCP2-dependent metabolic reprogramming in response to IL33. Upon tissue injury, acts downstream of IL33 signaling to drive differentiation of inflammation-resolving alternatively activated macrophages.

Subcellular Localization: Nucleus .

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer-composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.