

Recombinant Mouse IL27B Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP52026

Size: 10ug, 100ug

Protein Name: Interleukin-27 subunit beta (IL-27 subunit beta) (IL-27B) (Epstein-Barr virus-induced gene 3 protein homolog)

Synonym: Ebi3, Il27b

UniProt ID: O35228

UniProt Link: www.uniprot.org/uniprotkb/O35228/entry

Species Source: Mouse

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 228 AA

Molecular Weight: 25 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MSKLLFLSLALWASRSPGYTETALVALSQPRVQCHASRYPVAVDCSWTPLQAPNSTRSTSFIATYRL
GVATQQSQQPCLQRSPQASRCTIPDVHLFSTVPYMLNVTAVHPGGASSLLAFVAERIIKPDPEGV
RLRTAGQRLQVLWHPPASWPFDPDIFSLKYRLRYRRRGASHFRQVGPIEATTFTLRNSKPHAKYCIQV
SAQDLTDYGKPSDWSLPGQVESAPHKP

3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.

Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant Interleukin-27 subunit beta (IL-27 subunit beta) (IL-27B) (Epstein-Barr virus-induced gene 3 protein homolog) produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: Associates with IL27 to form the IL-27 interleukin, a heterodimeric cytokine which functions in innate immunity. IL-27 has pro- and anti-inflammatory properties, that can regulate T-helper cell development, suppress T-cell proliferation, stimulate cytotoxic T-cell activity, induce isotype switching in B-cells, and that has diverse effects on innate immune cells. Among its target cells are CD4 T-helper cells which can differentiate in type 1 effector cells (TH1), type 2 effector cells (TH2) and IL17 producing helper T-cells (TH17). It drives rapid clonal expansion of naive but not memory CD4 T-cells. It also strongly synergizes with IL-12 to trigger interferon-gamma/IFN-gamma production of naive CD4 T-cells, binds to the cytokine receptor WSX-1/TCR. Another important role of IL-27 is its antitumor activity as well as its antiangiogenic activity with activation of production of antiangiogenic chemokines.

Subcellular Localization: Secreted .

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer-composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.