

Recombinant Mouse NGF Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP51208

Size: 10ug, 100ug

Protein Name: Beta-nerve growth factor (Beta-NGF)

Synonym: Ngf, Ngfb

UniProt ID: P01139

UniProt Link: www.uniprot.org/uniprotkb/P01139/entry

Species Source: Mouse

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 241 AA

Molecular Weight: 27 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MSMLFYTLITAFILIGVQAEPYTDSNVPEGDSVPEAHWTKLQHSLDTALRRARSAPTAPIAARVTGQTRNITVDPRLFKKRRLLHSPRVLFSTQPPPTSSDTLDLDFQAHGTIPFNRTHRSKRSSTHPVFHMGFEFSCDSVSVWVGDKTTATDIKGKEVTVLAEVNINNSVFRQYFFETKCRASNPVESGCRGIDSKHWNSYCTTHTFVKALTTDEKQAAWRFIRIDTACVCVLSRKATRRG

3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.

Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant Beta-nerve growth factor (Beta-NGF) produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: Nerve growth factor is important for the development and maintenance of the sympathetic and sensory nervous systems (PubMed:20036257). Extracellular ligand for the NTRK1 and NGFR receptors, activates cellular signaling cascades to regulate neuronal proliferation, differentiation and survival (PubMed:22649032). The immature NGF precursor (proNGF) functions as a ligand for the heterodimeric receptor formed by SORCS2 and NGFR, and activates cellular signaling cascades that lead to inactivation of RAC1 and/or RAC2, reorganization of the actin cytoskeleton and neuronal growth cone collapse (PubMed:22155786). In contrast to mature NGF, the precursor form (proNGF) promotes neuronal apoptosis (in vitro) (PubMed:20036257). Inhibits metalloproteinase-dependent proteolysis of platelet glycoprotein VI (By similarity). Binds lysophosphatidylinositol and lysophosphatidylserine between the two chains of the homodimer (PubMed:22649032, PubMed:26144237). The lipid-bound form promotes histamine release from mast cells, contrary to the lipid-free form (PubMed:22649032).

Subcellular Localization: Secreted . Endosome lumen . Note=ProNGF is endocytosed after binding to the cell surface receptor formed by SORT1 and NGFR.

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer-composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.