

Recombinant Human TLE5 Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP50968

Size: 10ug, 100ug

Protein Name: TLE family member 5 (Amino-terminal enhancer of split) (Amino enhancer of split) (Gp130-associated protein GAM) (Grg-5) (Groucho-related protein 5) (Protein ESP1) (Protein GRG) (TLE family member 5, transcriptional modulator)

Synonym: TLE5, AES, GRG, GRG5

UniProt ID: Q08117

UniProt Link: www.uniprot.org/uniprotkb/Q08117/entry

Species Source: Human

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 197 AA

Molecular Weight: 22 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MMFPQSRHSGSSHLPQQLKFTTSDSCDRIKDEFQLLQAQYHSLKLECDKLASEKSEMQRHYVMYY
EMSYGLNIEMHKQAEIVKRLNGICAQVLPYLSQEHHQQVVLGAIERAKQVTAPELNSIIRQQLQAHQLS
QLQALALPLTLPVGLQPPSLPAVSAGTGLLSLSALGSQAHLISKEDKNGHDGDTHQEDDGEKSD

3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.

Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant TLE family member 5 (Amino-terminal enhancer of split) (Amino enhancer of split) (Gp130-associated protein GAM) (Grg-5) (Groucho-related protein 5) (Protein ESP1) (Protein GRG) (TLE family member 5, transcriptional modulator) produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: Transcriptional corepressor. Acts as a dominant repressor towards other family members. Inhibits NF-kappa-B-regulated gene expression. May be required for the initiation and maintenance of the differentiated state. Essential for the transcriptional repressor activity of SIX3 during retina and lens development.

Subcellular Localization: Nucleus .

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer-composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.