

Recombinant Mouse HCST Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP51947

Size: 10ug, 100ug

Protein Name: Hematopoietic cell signal transducer (DNAX-activation protein 10)
(Membrane protein DAP10) (Transmembrane adapter protein KAP10)

Synonym: Hcst, Dap10, Kap10

UniProt ID: Q9QUJ0

UniProt Link: www.uniprot.org/uniprotkb/Q9QUJ0/entry

Species Source: Mouse

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 79 AA

Molecular Weight: 8 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MDPPGYLLFLLLLPVAASQTSAGSCSGCGLSLPLLAGLVAAADAVMSLLIVGVVFVCMRPHGRPAQE
DGRVYINMPGRG

3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.

Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant Hematopoietic cell signal transducer (DNAX-activation protein 10) (Membrane protein DAP10) (Transmembrane adapter protein KAP10) produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: Transmembrane adapter protein which associates with KLRK1 to form an activation receptor KLRK1-HCST in lymphoid and myeloid cells; this receptor plays a major role in triggering cytotoxicity against target cells expressing cell surface ligands such as MHC class I chain-related MICA and MICB, and UL16-binding proteins (ULBPs); these ligands are up-regulated by stress conditions and pathological state such as viral infection and tumor transformation. Functions as a docking site for PI3-kinase PIK3R1 and GRB2. Interaction of ULBPs with KLRK1-HCST triggers calcium mobilization and activation of the PIK3R1, MAP2K/ERK, and JAK2/STAT5 signaling pathways. Both PIK3R1 and GRB2 are required for full KLRK1-HCST-mediated activation and ultimate killing of target cells. In NK cells, KLRK1-HCST signaling directly induces cytotoxicity and enhances cytokine production initiated via DAP12/TYROBP-associated receptors. In T-cells, it provides primarily costimulation for TCR-induced signals. KLRK1-HCST receptor plays a role in immune surveillance against tumors and is required for cytolysis of tumors cells; indeed, melanoma cells that do not express KLRK1 ligands escape from immune surveillance mediated by NK cells.

Subcellular Localization: Membrane ; Single-pass type I membrane protein .

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer-composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.