

Recombinant Rat CEAM1 Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP52284

Size: 10ug, 100ug

Protein Name: Cell adhesion molecule CEACAM1 (ATP-dependent taurocolate-carrier protein) (Carcinoembryonic antigen-related cell adhesion molecule 1) (CEA cell adhesion molecule 1) (Cell-CAM 105) (C-CAM 105) (Ecto-ATPase) (GP110) (pp120) (CD antigen CD66a)

Synonym: Ceacam1

UniProt ID: P16573

UniProt Link: www.uniprot.org/uniprotkb/P16573/entry

Species Source: Rat

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 519 AA

Molecular Weight: 57 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MELASARLLRGQIPWRGLLLTASLLTYWSPLTTAQVTVDVAPPNVVEEKSVLLLAHNLPQEFQVFYW
YKGTTLNPDSEIARYIRSDNMSKTGPAYSGRETIYSNGSLFFQNVNKTDERAYTLVFDQQFNPIQTS
VQFRVYPALQKPNVTGNNSNPMEGEPFVSLMCEPYTNNTSYLWSRNGESLSEGDRVTFSEGNRTL
TLLNVRRTDKGYECEARNPATFNRSDPFNLDVIYGPDAVISPPDIYLHQGSNLNLSCHADSNPPA
QYFWLINEKLQTSSQELFISNITTNSGTACFVNNTVTGLSRTTVKNITVFEPVTQPSIQITNTTVKEL
GSVTLTCSKDTGVSVRWLFNSQSLQLTDRMTLSQDNSTLRIDPIKREDAGDYQCEISNPVSFRISHP
IKLDVIPDPTQGNSGLSEGAIAIGVIGSVAGVALIAALAYFLYSRKTGGGSDHRDLTEHKPSTSSHNLG
PSDDSPNKVDDVSYSVLNFNAQQSKRPTSASSSPTETVYSVVKKK

3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.



Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant Cell adhesion molecule CEACAM1 (ATP-dependent taurocholate-carrier protein) (Carcinoembryonic antigen-related cell adhesion molecule 1) (CEA cell adhesion molecule 1) (Cell-CAM 105) (C-CAM 105) (Ecto-ATPase) (GP110) (pp120) (CD antigen CD66a) produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: [Isoform 1]: Cell adhesion protein that mediates homophilic cell adhesion in a calcium-independent manner (PubMed:2373740, PubMed:8454589). Plays a role as coinhibitory receptor in immune response, insulin action and also functions as an activator during angiogenesis (PubMed:11850617). Its coinhibitory receptor function is phosphorylation- and PTPN6 -dependent, which in turn, suppress signal transduction of associated receptors by dephosphorylation of their downstream effectors (By similarity). Plays a role in immune response, of T-cells, natural killer (NK) and neutrophils (By similarity). Upon TCR/CD3 complex stimulation, inhibits TCR-mediated cytotoxicity by blocking granule exocytosis by mediating homophilic binding to adjacent cells, allowing interaction with and phosphorylation by LCK and interaction with the TCR/CD3 complex which recruits PTPN6 resulting in dephosphorylation of CD247 and ZAP70 (By similarity). Also inhibits T-cell proliferation and cytokine production through inhibition of JNK cascade and plays a crucial role in regulating autoimmunity and anti-tumor immunity by inhibiting T-cell through its interaction with HAVCR2 (By similarity). Upon natural killer (NK) cells activation, inhibit KLRK1-mediated cytotoxicity of CEACAM1-bearing tumor cells by trans-homophilic interactions with CEACAM1 on the target cell and lead to cis-interaction between CEACAM1 and KLRK1, allowing PTPN6 recruitment and then VAV1 dephosphorylation (By similarity). Upon neutrophils activation negatively regulates IL1B production by recruiting PTPN6 to a SYK-TLR4-CEACAM1 complex, that dephosphorylates SYK, reducing the production of reactive oxygen species (ROS) and lysosome disruption, which in turn, reduces the activity of the inflammasome (By similarity). Down-regulates neutrophil production by acting as a coinhibitory receptor for CSF3R by down-regulating the CSF3R-STAT3 pathway through recruitment of PTPN6 that dephosphorylates CSF3R (By similarity). Also regulates insulin action by promoting INS clearance and regulating lipogenesis in liver through regulating insulin signaling (PubMed:11850617). Upon INS stimulation, undergoes phosphorylation by INSR leading to INS clearance by increasing receptor-mediated insulin endocytosis (PubMed:7592607, PubMed:9712832). This internalization promotes interaction with FASN leading to receptor-mediated insulin degradation and to reduction of FASN activity leading to negative regulation of fatty acid synthesis (PubMed:16054098, PubMed:7592607). INSR-mediated phosphorylation also provokes a down-regulation of cell proliferation through SHC1 interaction resulting in decrease coupling of SHC1 to the MAPK3/ERK1-MAPK1/ERK2 and phosphatidylinositol 3-kinase pathways (PubMed:11694516). Functions as activator in angiogenesis by promoting blood vessel remodeling through endothelial cell differentiation and migration and in arteriogenesis by increasing the number of collateral arteries and collateral vessel calibers after ischemia (By similarity). Also regulates vascular permeability through the VEGFR2 signaling pathway resulting in control of nitric oxide production (By similarity). Down-regulates cell growth in response to EGF through its interaction with SHC1 that mediates interaction with EGFR resulting in decrease coupling of SHC1 to the MAPK3/ERK1-MAPK1/ERK2 pathway (PubMed:15467833). Negatively regulates platelet aggregation by decreasing platelet adhesion on type I collagen through the GPVI-FcRgamma complex (By similarity). Inhibits cell migration and cell scattering through interaction with FLNA; interferes with the interaction of FLNA with RALA (By similarity). Mediates bile acid transport activity

in a phosphorylation dependent manner (PubMed:7518458). Negatively regulates osteoclastogenesis (By similarity).; [Isoform 2]: Cell adhesion proteins that mediates homophilic cell adhesion in a calcium-independent manner (PubMed:7774714, PubMed:8536699). Promotes populations of T-cells regulating IgA production and secretion associated with control of the commensal microbiota and resistance to enteropathogens (By similarity).

Subcellular Localization: [Isoform 1]: Cell membrane ; Single-pass type I membrane protein. Lateral cell membrane . Apical cell membrane . Basal cell membrane . Cell junction . Cell junction, adherens junction . Note=Canalicular domain of hepatocyte plasma membranes (PubMed:8513803). Found as a mixture of monomer, dimer and oligomer in the plasma membrane. Occurs predominantly as cis-dimers and/or small cis-oligomers in the cell junction regions (PubMed:19948503). Found as dimer in the solution (PubMed:9003371). Predominantly localized to the lateral cell membranes (PubMed:7774714).; [Isoform 2]: Cell membrane ; Single-pass type I membrane protein. Lateral cell membrane . Apical cell membrane . Basal cell membrane . Cell junction . Cell junction, adherens junction . Cytoplasmic vesicle, secretory vesicle . Note=Predominantly localized to the lateral cell membranes (PubMed:7774714). Found as a mixture of monomer, dimer and oligomer in the plasma membrane. Occurs predominantly as cis-dimers and/or small cis-oligomers in the cell junction regions (PubMed:19948503). Co-localizes with ANXA2 in secretory vesicles and with S100A10/p11 at the plasma membrane (By similarity).; Cell projection, microvillus membrane ; Single-pass type I membrane protein . Apical cell membrane ; Single-pass type I membrane protein . Note=Localized to the apical glycocalyx surface (By similarity). Colocalizes with CEACAM20 at the apical brush border of intestinal cells (By similarity).

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer-composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.