

Recombinant Mouse IFNG Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP51214

Size: 10ug, 100ug

Protein Name: Interferon gamma (IFN-gamma)

Synonym: Ifng

UniProt ID: P01580

UniProt Link: www.uniprot.org/uniprotkb/P01580/entry

Species Source: Mouse

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 155 AA

Molecular Weight: 18 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MNATHCILALQLFLMAVSGCYCHGTVIESLES LN NYFNSSGIDVEEKSLFLDIWRNWQKDGMKILQS
QIISFYLR LFEVLKDNQAISNNISVIESHLITTF SNSKAKKDAFMSIAKFEVNNPQVQRQAFNELIRVVH
QLLPESLRKRKR SRC

3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.

Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant Interferon gamma (IFN-gamma) produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: Type II interferon produced by immune cells such as T-cells and NK cells that plays crucial roles in antimicrobial, antiviral, and antitumor responses by activating effector immune cells and enhancing antigen presentation (PubMed:11585387, PubMed:8456301). Primarily signals through the JAK-STAT pathway after interaction with its receptor IFNGR1 to affect gene regulation. Upon IFNG binding, IFNGR1 intracellular domain opens out to allow association of downstream signaling components JAK2, JAK1 and STAT1, leading to STAT1 activation, nuclear translocation and transcription of IFNG-regulated genes. Many of the induced genes are transcription factors such as IRF1 that are able to further drive regulation of a next wave of transcription. Plays a role in class I antigen presentation pathway by inducing a replacement of catalytic proteasome subunits with immunoproteasome subunits. In turn, increases the quantity, quality, and repertoire of peptides for class I MHC loading. Increases the efficiency of peptide generation also by inducing the expression of activator PA28 that associates with the proteasome and alters its proteolytic cleavage preference. Up-regulates as well MHC II complexes on the cell surface by promoting expression of several key molecules such as cathepsins B/CTSB, H/CTSH, and L/CTSL (By similarity). Participates in the regulation of hematopoietic stem cells during development and under homeostatic conditions by affecting their development, quiescence, and differentiation (PubMed:20535209, PubMed:25078851).

Subcellular Localization: Secreted .

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer-composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.