

Recombinant Mouse SLAF1 Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP51950

Size: 10ug, 100ug

Protein Name: Signaling lymphocytic activation molecule (SLAM family member 1) (CD antigen CD150)

Synonym: Slamf1, Slam

UniProt ID: Q9QUM4

UniProt Link: www.uniprot.org/uniprotkb/Q9QUM4/entry

Species Source: Mouse

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 343 AA

Molecular Weight: 38 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MDPKGSLSWRILLFLSLAFELSYGTGGGVMDCPVILQKLGQDTWLPLTNEHQINKSVNKSVRILVTM
ATSPGSKSNKKIVSFDLSKGSYPDHLEDGYHFQSKNLSLKILGNRRESEGWYLVSVVEENVSVQQFC
KQLKLYEQVSPPEIKVLNKTQENENGTCSLLLACTVKKGDHVTYSWSDEAGTHLLSRANRSHLLHITL
SNQHQDSIYNCTASNPVSSISRTFNLSSQACKQESSSESSPWMQYTLVPLGVVIFILVFTAIIMMKRQ
GKSNHCQPPVEEKSLTIYAQVQKSGPQEKKLHDALTDQDPCTTIYVAATEPAPESVQEPNPTTVYAS
VTLPEs

3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.

Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions



Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant Signaling lymphocytic activation molecule (SLAM family member 1) (CD antigen CD150) produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: Self-ligand receptor of the signaling lymphocytic activation molecule (SLAM) family. SLAM receptors triggered by homo- or heterotypic cell-cell interactions are modulating the activation and differentiation of a wide variety of immune cells and thus are involved in the regulation and interconnection of both innate and adaptive immune response. Activities are controlled by presence or absence of small cytoplasmic adapter proteins, SH2D1A/SAP and/or SH2D1B/EAT-2. SLAMF1-induced signal-transduction events in T-lymphocytes are different from those in B-cells. Two modes of SLAMF1 signaling seem to exist: one depending on SH2D1A (and perhaps SH2D1B) and another in which protein-tyrosine phosphatase 2C (PTPN11)-dependent signal transduction operates. Initially it has been proposed that association with SH2D1A prevents binding to inhibitory effectors including INPP5D/SHIP1 and PTPN11/SHP-2 (By similarity). However, signaling is also regulated by SH2D1A which can simultaneously interact with and recruit FYN which subsequently phosphorylates and activates SLAMF1 (By similarity). Mediates IL-2-independent proliferation of activated T-cells during immune responses and induces IFN-gamma production (PubMed:12351401, PubMed:9126961). Downstreaming signaling involves INPP5D, DOK1 and DOK2 leading to inhibited IFN-gamma production in T-cells, and PRKCQ, BCL10 and NFKB1 leading to increased T-cell activation and Th2 cytokine production (PubMed:11477403, PubMed:15539155, PubMed:16847311). Promotes T-cell receptor-induced IL-4 secretion by CD4(+) cells (PubMed:15123745). Inhibits antigen receptor-mediated production of IFN-gamma, but not IL-2, in CD4(-)/CD8(-) T-cells (PubMed:11477403). Required for IL-4 production by germinal centers T follicular helper (T(Fh)) cells (PubMed:20525889). May inhibit CD40-induced signal transduction in monocyte-derived dendritic cells (By similarity). May play a role in allergic responses and may regulate allergen-induced Th2 cytokine and Th1 cytokine secretion (PubMed:16528012). In conjunction with SLAMF6 controls the transition between positive selection and the subsequent expansion and differentiation of the thymocytic natural killer T (NKT) cell lineage (PubMed:18031695). Involved in the peripheral differentiation of indifferent natural killer T (iNKT) cells toward a regulatory NKT2 type (PubMed:18606638). In macrophages involved in down-regulation of IL-12, TNF and nitric oxide in response to lipopolysaccharide (LPS) (PubMed:15123745). In B-cells activates the ERK signaling pathway independently of SH2D1A but implicating both, SYK and INPP5D, and activates Akt signaling dependent on SYK and SH2D1A (PubMed:15315965). In conjunction with CD84/SLAMF5 and SLAMF6 may be a negative regulator of the humoral immune response (PubMed:25926831).; (Microbial infection) Involved in innate immune response against Gram-negative bacteria in macrophages; probably recognizes OmpC and/or OmpF on the bacterial surface, regulates phagosome maturation and recruitment of the PI3K complex II (PI3KC3-C2) leading to accumulated of PtdIns(3)P and NOX2 activity in the phagosomes (PubMed:20818396, PubMed:22493499).

Subcellular Localization: Cell membrane ; Single-pass type I membrane protein. Note=Present on the surface of B-cells and T-cells. Located at the plasma membrane contacts between neighboring T cells.

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer-composition requirements at the time of order placement.



Enlibio Wuhan Enlibio Biotech Co.,Ltd

- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.