

Recombinant Mouse IRF1 Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP51283

Size: 10ug, 100ug

Protein Name: Interferon regulatory factor 1 (IRF-1)

Synonym: Irf1

UniProt ID: P15314

UniProt Link: www.uniprot.org/uniprotkb/P15314/entry

Species Source: Mouse

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 329 AA

Molecular Weight: 37 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MPITRMRMRPWLEMQINSNQIPGLIWINKEEMIFQIPWKHAAKHGWDINKDACLFERSWAIHTGRYKA
GEKEPDPKTKWANKFRCAMNSLPDIEEVKDQSRNKGSSAVRVYRMLPPLTRNQRKERKSKSSRDTK
SKTKRKLCDGVSPDTFSDGLSSSTLPDDHSSYTTQGYLGQDLDMERDITPALSPCVSSSLSEWHM
QMDIIPDSTTDLYNLQVSPMPSTSEAATDEDEEGKIAEDLMKLFEQSEWQPTHIDGKGYYLLNEPGTQ
LSSVYGDFSCKEEPEIDSPRGDIGIGIQHVFTMKNMDSIMWMDSLLGNSVRLPPSIQAIPCAP

3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.

Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions



Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant Interferon regulatory factor 1 (IRF-1) produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: Transcriptional regulator which displays a remarkable functional diversity in the regulation of cellular responses (PubMed:11244049, PubMed:11846971, PubMed:11846974, PubMed:16932750, PubMed:20049431, PubMed:25774715). Regulates transcription of IFN and IFN-inducible genes, host response to viral and bacterial infections, regulation of many genes expressed during hematopoiesis, inflammation, immune responses and cell proliferation and differentiation, regulation of the cell cycle and induction of growth arrest and programmed cell death following DNA damage (PubMed:11244049, PubMed:11846971, PubMed:11846974, PubMed:16932750, PubMed:20049431, PubMed:2208287, PubMed:3409321, PubMed:8402903). Stimulates both innate and acquired immune responses through the activation of specific target genes and can act as a transcriptional activator and repressor regulating target genes by binding to an interferon-stimulated response element (ISRE) in their promoters (PubMed:11244049, PubMed:11846971, PubMed:11846974, PubMed:16932750, PubMed:20049431). Has an essential role in IFN γ -dependent immunity to mycobacteria (By similarity). Binds to a consensus sequence in gene promoters (By similarity). Its target genes for transcriptional activation activity are: genes involved in anti-viral response, such as IFN- α /beta, RIGI, TNFSF10/TRAIL, ZBP1, OAS1/2, PIAS1/GBP, EIF2AK2/PKR and RSAD2/viperin; antibacterial response, such as GBP2, GBP5, IRGB10 and NOS2/INOS; anti-proliferative response, such as p53/TP53, LOX and CDKN1A; apoptosis, such as BBC3/PUMA, CASP1, CASP7 and CASP8; immune response, such as IL7, IL12A/B and IL15, PTGS2/COX2 and CYBB; DNA damage responses and DNA repair, such as POLQ/POLH; MHC class I expression, such as TAP1, PSMB9/LMP2, PSME1/PA28A, PSME2/PA28B and B2M and MHC class II expression, such as CIITA; metabolic enzymes, such as ACOD1/IRG1 (PubMed:12387893, PubMed:17018642, PubMed:18955028, PubMed:20308629, PubMed:21909274, PubMed:25774715, PubMed:27693356, PubMed:29321274, PubMed:30635240). Represses genes involved in anti-proliferative response, such as BIRC5/survivin, CCNB1, CCNE1, CDK1, CDK2 and CDK4 and in immune response, such as FOXP3, IL4, ANXA2 and TLR4 (PubMed:18641303). Stimulates p53/TP53-dependent transcription through enhanced recruitment of EP300 leading to increased acetylation of p53/TP53 (By similarity). Plays an important role in immune response directly affecting NK maturation and activity, macrophage production of IL12, Th1 development and maturation of CD8 $^{+}$ T-cells (PubMed:11244049, PubMed:11846971, PubMed:11846974, PubMed:16932750, PubMed:20049431). Also implicated in the differentiation and maturation of dendritic cells and in the suppression of regulatory T (Treg) cells development (PubMed:11244049, PubMed:11846971, PubMed:11846974, PubMed:16932750, PubMed:20049431). Acts as a tumor suppressor and plays a role not only in antagonism of tumor cell growth but also in stimulating an immune response against tumor cells (PubMed:11244049, PubMed:11846971, PubMed:11846974, PubMed:16932750, PubMed:20049431).

Subcellular Localization: Nucleus . Cytoplasm . Note=MYD88-associated IRF1 migrates into the nucleus more efficiently than non-MYD88-associated IRF1.

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer-composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial

manufacturing.