

# Recombinant Mouse LOX5 Protein

**For Research Use Only**

## 1. Basic Product Information

Catalog No.: REP51444

Size: 10ug, 100ug

Protein Name: Polyunsaturated fatty acid 5-lipoxygenase (EC 1.13.11.-) (Arachidonate 5-lipoxygenase) (5-LO) (5-lipoxygenase) (EC 1.13.11.34)

Synonym: Alox5

UniProt ID: P48999

UniProt Link: [www.uniprot.org/uniprotkb/P48999/entry](http://www.uniprot.org/uniprotkb/P48999/entry)

Species Source: Mouse

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 674 AA

Molecular Weight: 78 KD

Protein Tag: C-terminal His6

## 2. Amino Acid Sequence

MPSYTVTVATGSQWFAGTDDYIYLSLIGSAGCSEKHLLDKAFYNDFERGAVDSYDVTVDDEELGEIYL  
VKIEKRKYWLHDDWYLKYITLKTTPHGDYIEFPCYRWITGEGEIVLRDGRAKLARDDQIHILKQHRRKEL  
EARQKQYRWMEWNPGFPLSIDAKCHKDLPRDIQFDSEKGVDFVLNYSKAMENLFINRFMHMFQSS  
WHDFADFEKIFVKISNTISERVKNHWQEDLMFGYQFLNGCNPVLIKRCTALPPKLPVTTEMVECSLER  
QLSLEQEVQEGNIFVDYELLDGIDANKTDPCTHQFLAAPICLLYKNLANKIVPIAIQLNQTPGESNPIFL  
PTDSKYDWLLAKIWVRSSDFHVHQTITHLLRTHLVSEVFGIAMYRQLPAVHPLFKLLVAHVRFTIAINT  
KAREQLICEYGLFDKANATGGGGHVQMVRVQDLTYSSLCFPEAIKARGMDSTEDIPFYFYRDDG  
LLVWEAIQSFTMEVVSIIYENDQVVEEDQELQDFVKDYYVYGMRGKKASGFPSIKSREKLSEYLT  
VIFTASAQHA AVNFGQYDWCSWIPNAPPTMRAPPTAKGVVTIEQIVDTLPDRGRSCWHLGAVWAL  
SQFQENELFLGMYPEEHFIEKPVKEAMIRFRKNLEAIVSVIAERNKNKKLPYLYLSPDRIPNSVAI

## 3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.



Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

## 4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

## 5. Product Description

This product is recombinant Polyunsaturated fatty acid 5-lipoxygenase (EC 1.13.11.-) (Arachidonate 5-lipoxygenase) (5-LO) (5-lipoxygenase) (EC 1.13.11.34) produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

**Protein Function:** Catalyzes the oxygenation of arachidonate to 5-hydroperoxyeicosatetraenoate (5-HPETE) followed by the dehydration to 5,6-epoxyeicosatetraenoate (Leukotriene A<sub>4</sub>/LTA<sub>4</sub>), the first two steps in the biosynthesis of leukotrienes, which are potent mediators of inflammation (PubMed:23246375, PubMed:31642348, PubMed:7629107, PubMed:7809134, PubMed:7969451). Also catalyzes the oxygenation of arachidonic acid into 8-hydroperoxyicosatetraenoic acid (8-HPETE) and 12-hydroperoxyicosatetraenoic acid (12-HPETE) (PubMed:23246375). Displays lipoxin synthase activity being able to convert (15S)-HETE into a conjugate tetraene (By similarity). Although arachidonate is the preferred substrate, this enzyme can also metabolize oxidized fatty acids derived from arachidonate such as (15S)-HETE, eicosapentaenoate (EPA) such as (18R)- and (18S)-HEPE or docosahexaenoate (DHA) which lead to the formation of specialized pro-resolving mediators (SPM) lipoxin and resolvins E and D respectively, therefore it participates in anti-inflammatory responses (PubMed:31642348). Oxidation of DHA directly inhibits endothelial cell proliferation and sprouting angiogenesis via peroxisome proliferator-activated receptor gamma (PPARgamma) (PubMed:21307302). It does not catalyze the oxygenation of linoleic acid and does not convert (5S)-HETE to lipoxin isomers (PubMed:31642348). In addition to inflammatory processes, participates in dendritic cell migration, wound healing through an antioxidant mechanism based on heme oxygenase-1 (HO-1) regulation expression, monocyte adhesion to the endothelium via ITGAM expression on monocytes (PubMed:17392829, PubMed:23720274, PubMed:24226420, PubMed:28965882). Moreover, it helps establish an adaptive humoral immunity by regulating primary resting B cells and follicular helper T cells and participates in the CD40-induced production of reactive oxygen species (ROS) after CD40 ligation in B cells through interaction with PIK3R1 that bridges ALOX5 with CD40 (PubMed:21224059). May also play a role in glucose homeostasis, regulation of insulin secretion and palmitic acid-induced insulin resistance via AMPK (PubMed:18421434, PubMed:28694473). Can regulate bone mineralization and fat cell differentiation increases in induced pluripotent stem cells (PubMed:24906289).

**Subcellular Localization:** Cytoplasm . Nucleus matrix . Nucleus membrane ; Peripheral membrane protein . Cytoplasm, perinuclear region . Cytoplasm, cytosol . Nucleus envelope . Nucleus intermembrane space . Note=Shuttles between cytoplasm and nucleus. Found exclusively in the nucleus, when phosphorylated on Ser-272. Calcium binding promotes translocation from the cytosol and the nuclear matrix to the nuclear envelope and membrane association.

## 6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer-composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial

manufacturing.