

Recombinant Mouse OST48 Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP52126

Size: 10ug, 100ug

Protein Name: Dolichyl-diphosphooligosaccharide--protein glycosyltransferase 48 kDa subunit (DDOST 48 kDa subunit) (Oligosaccharyl transferase 48 kDa subunit)

Synonym: Ddost

UniProt ID: O54734

UniProt Link: www.uniprot.org/uniprotkb/O54734/entry

Species Source: Mouse

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 441 AA

Molecular Weight: 49 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MKMDPRLAVRAWPLCGLLLAVALGCVCASGPRTLVLDDNLNVRDTHSLFFRSLKDRGFELTFKTADD
PSLSLIKYGEFLYDNLIIIFSPSVEDFGGNINVETISAFIDGGGSLVAASSDIGDPLRELGSECGIEFDEE
KTAVIDHHNYDVSDLGQHTLIVADTENLLKAPTIVGKSSLNPILFRGVGMVADPDNPLVLDILTGSSTS
YSFFPDKPITQYPHAVGRNTLLIAGLQARNNARVIFSGSLDFFSDAFFNSAVQKATPGAQRYSTGN
YELAVALSRWVFKEEGVLRVGPVSHHRVGEMAPPNAYTVTDLVEYSIIIEQLSNGKWVPFDGDDIQL
EFVRIDPFVRTFLKRKGGKYSVQFKLPDVYGVFQFKVDYNRLGYTHLYSSTQVSVRPLQHTQYERFI
PSAYPPYASAFSMMAGLFIFSIVFLHMKEKEKSD

3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.

Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant Dolichyl-diphosphooligosaccharide--protein glycosyltransferase 48 kDa subunit (DDOST 48 kDa subunit) (Oligosaccharyl transferase 48 kDa subunit) produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: Subunit of the oligosaccharyl transferase (OST) complex that catalyzes the initial transfer of a defined glycan (Glc(3)Man(9)GlcNAc(2) in eukaryotes) from the lipid carrier dolichol-pyrophosphate to an asparagine residue within an Asn-X-Ser/Thr consensus motif in nascent polypeptide chains, the first step in protein N-glycosylation (By similarity). N-glycosylation occurs cotranslationally and the complex associates with the Sec61 complex at the channel-forming translocon complex that mediates protein translocation across the endoplasmic reticulum (ER). All subunits are required for a maximal enzyme activity (By similarity). Required for the assembly of both SST3A- and SS3B-containing OST complexes (By similarity).

Subcellular Localization: Endoplasmic reticulum membrane ; Single-pass type I membrane protein .

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer-composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.