

Recombinant Mouse TN13B Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP51981

Size: 10ug, 100ug

Protein Name: Tumor necrosis factor ligand superfamily member 13B (B-cell-activating factor) (BAFF) (CD antigen CD257) [Cleaved into: Tumor necrosis factor ligand superfamily member 13b, membrane form; Tumor necrosis factor ligand superfamily member 13b, soluble form]

Synonym: Tnfsf13b, Baff

UniProt ID: Q9WU72

UniProt Link: www.uniprot.org/uniprotkb/Q9WU72/entry

Species Source: Mouse

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 309 AA

Molecular Weight: 34 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MDESAKTLPPPCLCFCSEKGEDMKVGYDPITPQKEEGAWFGICRDGRLLAATLLLALLSSSFTAMSL
YQLAALQADLMNLRMELQSYRGSATPAAAGAPELTAGVKLLTPAAPRPHNSSRGHRNRRAFQGPE
ETEQDVDLSAPPAPCLPGCRHSQHDDNGMNLRNIIQDCLQLIADSDTPTIRKGTYTFVPWLLSFKRG
NALEEKENKIVVRQTGYFFIYSQVLYTDPIFAMGHVIQRKKVHVFGDELSLVTLFRCIQNMPKTLPNNS
CYSAGIARLEEGDEIQLAIPRENAQISRNGDDTFFGALKLL

3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.

Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant Tumor necrosis factor ligand superfamily member 13B (B-cell-activating factor) (BAFF) (CD antigen CD257) [Cleaved into: Tumor necrosis factor ligand superfamily member 13b, membrane form; Tumor necrosis factor ligand superfamily member 13b, soluble form] produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: Cytokine that binds to TNFRSF13B/TACI and TNFRSF17/BCMA. TNFSF13/APRIL binds to the same 2 receptors. Together, they form a 2 ligands -2 receptors pathway involved in the stimulation of B- and T-cell function and the regulation of humoral immunity. A third B-cell specific BAFF-receptor (BAFFR/BR3) promotes the survival of mature B-cells and the B-cell response.; Isoform 2 seems to inhibit isoform 1 secretion and bioactivity.

Subcellular Localization: Cell membrane; Single-pass type II membrane protein.; [Tumor necrosis factor ligand superfamily member 13b, soluble form]: Secreted.

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer-composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.