

Recombinant Mouse BCL10 Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP51997

Size: 10ug, 100ug

Protein Name: B-cell lymphoma/leukemia 10 (B-cell CLL/lymphoma 10) (Bcl-10) (CARD-containing molecule enhancing NF-kappa-B) (CARD-like apoptotic protein) (mCLAP) (CED-3/ICH-1 prodomain homologous E10-like regulator) (mCIPER) (Cellular homolog of vCARMEN) (cCARMEN) (Cellular-E10) (c-E10) (Mammalian CARD-containing adapter molecule E10) (mE10)

Synonym: Bcl10, Ciper, Clap

UniProt ID: Q9Z0H7

UniProt Link: www.uniprot.org/uniprotkb/Q9Z0H7/entry

Species Source: Mouse

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 233 AA

Molecular Weight: 26 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MEAPAPSLTEEDLTEVKKDALENLRVYLCEKIIAERHFDHLRAKKILSREDTEEISCR TSSRK RAGKLL
DYLQENPRGLDTLVESIRREKTQSFLIQKITDEV LKLRNIKLEHLKGLKCSSCEPFAAGATNNLSRCNS
DESNLSEKQRASTVMYHPEGESSTAPFFSMASSLNLPVLEVGR TENS SFSSATLPRPGDPGAPPLP
PDLRLEEGGSCGNSSEMFLPLRSRALS RQ

3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.

Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant B-cell lymphoma/leukemia 10 (B-cell CLL/lymphoma 10) (Bcl-10) (CARD-containing molecule enhancing NF-kappa-B) (CARD-like apoptotic protein) (mCLAP) (CED-3/ICH-1 prodomain homologous E10-like regulator) (mCIPER) (Cellular homolog of vCARMEN) (cCARMEN) (Cellular-E10) (c-E10) (Mammalian CARD-containing adapter molecule E10) (mE10) produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: Plays a key role in both adaptive and innate immune signaling by bridging CARD domain-containing proteins to immune activation (PubMed:22265677). Acts by channeling adaptive and innate immune signaling downstream of CARD domain-containing proteins CARD9, CARD11 and CARD14 to activate NF-kappa-B and MAP kinase p38 (MAPK11, MAPK12, MAPK13 and/or MAPK14) pathways which stimulate expression of genes encoding pro-inflammatory cytokines and chemokines (PubMed:22265677). Recruited by activated CARD domain-containing proteins: homooligomerized CARD domain-containing proteins form a nucleating helical template that recruits BCL10 via CARD-CARD interaction, thereby promoting polymerization of BCL10, subsequent recruitment of MALT1 and formation of a CBM complex (By similarity). This leads to activation of NF-kappa-B and MAP kinase p38 (MAPK11, MAPK12, MAPK13 and/or MAPK14) pathways which stimulate expression of genes encoding pro-inflammatory cytokines and chemokines (By similarity). Activated by CARD9 downstream of C-type lectin receptors; CARD9-mediated signals are essential for antifungal immunity (PubMed:22265677). Activated by CARD11 downstream of T-cell receptor (TCR) and B-cell receptor (BCR) (By similarity). Promotes apoptosis, pro-caspase-9 maturation and activation of NF-kappa-B via NIK and IKK (By similarity).

Subcellular Localization: Cytoplasm . Membrane raft . Note=Colocalized with DPP4 in membrane rafts.

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer-composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.