

Recombinant Human INGR1 Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP50554

Size: 10ug, 100ug

Protein Name: Interferon gamma receptor 1 (IFN-gamma receptor 1) (IFN-gamma-R1) (CDw119) (Interferon gamma receptor alpha-chain) (IFN-gamma-R-alpha) (CD antigen CD119)

Synonym: IFNGR1

UniProt ID: P15260

UniProt Link: www.uniprot.org/uniprotkb/P15260/entry

Species Source: Human

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 489 AA

Molecular Weight: 54 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MALLFLLPLVMQGVSR AEMGTADLGPSSVPTPTNVTIESYNMNP IYWEYQIMPQVPVFTVEVKNYG
VKNSEWIDACINISHHYCNISDHVGDPSNSLWVRVKARVGQKESAYAKSEEFVAVCRDGKIGPPKLDI
RKEEKQIMIDIFHPSVFNVDGEQEVDPETTCYIRVYNVYVRMNGSEIQYKILTQKEDDCDEIQCQL
AIPVSSLNSQYCVSAEGVLHVWGVTTTEKSKEVCITIFNSSIKGSLWIPVVAALLFLVLSLVFICFYIKKI
NPLKEKSII LPKSLISVVR SATLETKPESKYVSLITSYQPFSLKEVVC EEPLSPATVPGMHTEDNPGK
VEHTEELSSITEVVTTEENIPDVVPGSHLTPIERESSPLSSNQSEPGSIALNSYHSRNCSESDHSRN
GFDTDSSCLES HSSLSDSEFPNNGEIKTEGQELITVIKAPTSFGYDKPHVLVDLLVDDSGKESLIGY
RPTEDSKEFS

3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.

Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant Interferon gamma receptor 1 (IFN-gamma receptor 1) (IFN-gamma-R1) (CDw119) (Interferon gamma receptor alpha-chain) (IFN-gamma-R-alpha) (CD antigen CD119) produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: Receptor subunit for interferon gamma/INFG that plays crucial roles in antimicrobial, antiviral, and antitumor responses by activating effector immune cells and enhancing antigen presentation (PubMed:20015550). Associates with transmembrane accessory factor IFNGR2 to form a functional receptor (PubMed:10986460, PubMed:2971451, PubMed:7615558, PubMed:7617032, PubMed:7673114). Upon ligand binding, the intracellular domain of IFNGR1 opens out to allow association of downstream signaling components JAK1 and JAK2. In turn, activated JAK1 phosphorylates IFNGR1 to form a docking site for STAT1. Subsequent phosphorylation of STAT1 leads to dimerization, translocation to the nucleus, and stimulation of target gene transcription (PubMed:28883123). STAT3 can also be activated in a similar manner although activation seems weaker. IFNGR1 intracellular domain phosphorylation also provides a docking site for SOCS1 that regulates the JAK-STAT pathway by competing with STAT1 binding to IFNGR1 (By similarity).

Subcellular Localization: Cell membrane ; Single-pass type I membrane protein .

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer-composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.