

Recombinant Mouse ITK Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP51556

Size: 10ug, 100ug

Protein Name: Tyrosine-protein kinase ITK/TSK (EC 2.7.10.2) (IL-2-inducible T-cell kinase) (Kinase EMT) (Kinase TLK) (T-cell-specific kinase)

Synonym: Itk, Emt, Tlk, Tsk

UniProt ID: Q03526

UniProt Link: www.uniprot.org/uniprotkb/Q03526/entry

Species Source: Mouse

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 625 AA

Molecular Weight: 72 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MNNFILLEEQLIKKSQQKRRTSPSNFKVRFFVLTKASLAYFEDRHGKKRTLKGSIELSRIKCVEIVKSDI
SIPCHYKYPFQTLVYLQVVHDNYLLYVFAPDCESRQRWVLTKEETRNNNSLVSKYHPNFWMDGR
WRCCSQLEKPAVGCAPYDPSKNASKKPLPPTPEDNRRSFQEPEETLVIALYDYQTNDPQELALRCD
EEYYLLDSSEIHWWRVQDKNGHEGYAPSSYLVEKSPNNLETYEWYNKSISRDKAEKLLDGTGKEGA
FMVRDSRTPGTYSVFTKAIISENPCIKHYHIKETNDSPKRYVVAEKYVFDSIPLLIQYHQYNGGGLV
TRLRYPVCSWRQKAPVTAGLRYGKWVIQPSSELTQVQEGSGQFGLVHLGYWLNKDKVAIKTIQEGA
MSEEDFIEEAEVMMKLSHPKLVQLYGVCLQAPICLVFEFMEHGCLSDYLRSGRGLFAAETLLGMCL
DVCEGMAYLEKACVIHRDLAARNCLVGENQVIKVSDFGMTRFVLDDQYTSSTGTGKFPVKWASPEVF
SFSRYSSKSDVWSFGVLMWEVFSEGEKIPYENRSNSEVVEDISTGFRLYKPRLASCHVYQIMNHCWK
EKPEDRPPFSQLLSQLAEIAEAGL

3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.

Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant Tyrosine-protein kinase ITK/TSK (EC 2.7.10.2) (IL-2-inducible T-cell kinase) (Kinase EMT) (Kinase TLK) (T-cell-specific kinase) produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: Tyrosine kinase that plays an essential role in regulation of the adaptive immune response. Regulates the development, function and differentiation of conventional T-cells and nonconventional NKT-cells. When antigen presenting cells (APC) activate T-cell receptor (TCR), a series of phosphorylation lead to the recruitment of ITK to the cell membrane, in the vicinity of the stimulated TCR receptor, where it is phosphorylated by LCK. Phosphorylation leads to ITK autophosphorylation and full activation. Once activated, phosphorylates PLCG1, leading to the activation of this lipase and subsequent cleavage of its substrates. In turn, the endoplasmic reticulum releases calcium in the cytoplasm and the nuclear activator of activated T-cells (NFAT) translocates into the nucleus to perform its transcriptional duty. Phosphorylates 2 essential adapter proteins: the linker for activation of T-cells/LAT protein and LCP2. Then, a large number of signaling molecules such as VAV1 are recruited and ultimately lead to lymphokine production, T-cell proliferation and differentiation. Required for TCR-mediated calcium response in gamma-delta T-cells, may also be involved in the modulation of the transcriptomic signature in the Vgamma2-positive subset of immature gamma-delta T-cells (PubMed:23562159). Phosphorylates TBX21 at 'Tyr-525' and mediates its interaction with GATA3 (PubMed:15662016).

Subcellular Localization: Cytoplasm. Nucleus . Note=Localizes in the vicinity of cell surface receptors in the plasma membrane after receptor stimulation.

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer-composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.