

Recombinant Mouse TNF14 Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP51960

Size: 10ug, 100ug

Protein Name: Tumor necrosis factor ligand superfamily member 14 (CD antigen CD258)
[Cleaved into: Tumor necrosis factor ligand superfamily member 14, membrane form; Tumor necrosis factor ligand superfamily member 14, soluble form]

Synonym: Tnfsf14, Light

UniProt ID: Q9QYH9

UniProt Link: www.uniprot.org/uniprotkb/Q9QYH9/entry

Species Source: Mouse

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 239 AA

Molecular Weight: 26 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MESVVQPSVVFVDGQTDIPFRRLEQNHRRRRCGTVQVSLALVLLGAGLATQGWFLRLHQRLGDI
VAHLPGDGKGSGWEKLIQDQRSHQANPAAHLTGANASLIGIGGPLLWETRLGLAFLRGLTYHDGALVT
MEPGYYYVYSKVQLSGVGCPQGLANGLPITHGLYKRTSRYPKLELLVSRRSPCGRANSSRVWWD
SSFLGGVVHLEAGEEVVVRVPGNRLVRPRDGTRSYFGAFMV

3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.

Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant Tumor necrosis factor ligand superfamily member 14 (CD antigen CD258) [Cleaved into: Tumor necrosis factor ligand superfamily member 14, membrane form; Tumor necrosis factor ligand superfamily member 14, soluble form] produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: Cytokine that binds to TNFRSF3/LTBR. Activates NFκB and stimulates the proliferation of T-cells. Acts as a ligand for TNFRSF14/HVEM. Upon binding to TNFRSF14/HVEM, delivers costimulatory signals to T cells, leading to T cell proliferation and IFNG production (By similarity).

Subcellular Localization: Cell membrane ; Single-pass type II membrane protein .; [Tumor necrosis factor ligand superfamily member 14, soluble form]: Secreted .

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer-composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.