

Recombinant Mouse TLR6 Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP51891

Size: 10ug, 100ug

Protein Name: Toll-like receptor 6 (CD antigen CD286)

Synonym: Tlr6

UniProt ID: Q9EPW9

UniProt Link: www.uniprot.org/uniprotkb/Q9EPW9/entry

Species Source: Mouse

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 795 AA

Molecular Weight: 91 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MSQDRKPIVGSFHFVCALALIVGSMTSPFSNELESMVDYSNRNLTHVPKDLPPRTKALSLSQNSISELR
MPDISFLSELRVLRSLSHNRIRSLDFHVFLFNQDLEYLDVSHNRLQNISCCPMASLRHLDLSFNDFDVL
PVCKEFGNLTCLTFLGLSAAKFRQLDLLPVAHLHLSCILLDLVSYHIKGGETESLQIPNTTVLHLVFHPN
SLFSVQVNMSVNALGHLQLSNIKLNDENCQRLMTFLSELTRGPTLLNVTLQHIETTWKCSVKLFQFF
WPRPVEYLNINLTITERIDREEFTYSETALKSLMIEHVKNQVFLFSKEALYSVFAEMNIKMLSISDTPFI
HMCPPSPSSFTFLNFTQNVFTDSVFQGCSTLKRQLTLILQRNGLKNFFKVALMTKNMSSLETLDVS
LNSLNSHAYDRTCAWAESILVLNLSSNMLTGSVFRCLPPKVKVLDLHNNRIMSIPKDVTHLQALQELN
VASNSLTDLPGCGAFSSLSVLVIDHNSVSHPSSEDFQSCQNIRSLTAGNNPFQCTCEL RDFVKNIGW
VAREVVEGWPD SYRCDYPESSKGTALRDFHMSPLSCDTVLLTVTIGATMLVLAVTGAFLCLYFDLPW
YVRMLCQWTQTRHRARHIPLEELQRNLQFHAFVSYSSEHDSAWVKNE LLPNLEKDDIRVCLHERNFV
PGKSIVENIINFIEKSYKAIFVLSPHFIQSEWCHYELYFAHNNLFHEGSDNLILILLEPILQNNIPSRYHKL
RALMAQRTYLEWPTEKGKRGLFWANLRASFIMKLALVNEDDVKT

3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.



Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant Toll-like receptor 6 (CD antigen CD286) produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: Participates in the innate immune response to Gram-positive bacteria and fungi. Specifically recognizes diacylated and, to a lesser extent, triacylated lipopeptides (PubMed:19931471). In response to diacylated lipopeptides, forms the activation cluster TLR2:TLR6:CD14:CD36, this cluster triggers signaling from the cell surface and subsequently is targeted to the Golgi in a lipid-raft dependent pathway. Acts via MYD88 and TRAF6, leading to NF-kappa-B activation, cytokine secretion and the inflammatory response. Recognizes mycoplasmal macrophage-activating lipopeptide-2kD (MALP-2), soluble tuberculosis factor (STF), phenol-soluble modulin (PSM) and B.burgdorferi outer surface protein A lipoprotein (OspA-L) cooperatively with TLR2. In complex with TLR4, promotes sterile inflammation in monocytes/macrophages in response to oxidized low-density lipoprotein (oxLDL) or amyloid-beta 42. In this context, the initial signal is provided by oxLDL- or amyloid-beta 42-binding to CD36. This event induces the formation of a heterodimer of TLR4 and TLR6, which is rapidly internalized and triggers inflammatory response, leading to the NF-kappa-B-dependent production of CXCL1, CXCL2 and CCL9 cytokines, via MYD88 signaling pathway, and CCL5 cytokine, via TICAM1 signaling pathway, as well as IL1B secretion (PubMed:20037584, PubMed:23812099).

Subcellular Localization: Cell membrane ; Single-pass type I membrane protein . Cytoplasmic vesicle, phagosome membrane ; Single-pass type I membrane protein . Membrane raft . Golgi apparatus . Note=Upon complex formation with CD36 and TLR4, internalized through dynamin-dependent endocytosis. Does not reside in lipid rafts before stimulation but accumulates increasingly in the raft upon the presence of the microbial ligand. In response to diacylated lipoproteins, TLR2:TLR6 heterodimers are recruited in lipid rafts, this recruitment determine the intracellular targeting to the Golgi apparatus.

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer-composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.