

Recombinant Mouse ANM5 Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP51769

Size: 10ug, 100ug

Protein Name: Protein arginine N-methyltransferase 5 (Prmt5) (EC 2.1.1.320) (Histone-arginine N-methyltransferase PRMT5) (Jak-binding protein 1) (Shk1 kinase-binding protein 1 homolog) (SKB1 homolog)

Synonym: Prmt5, Jbp1, Skb1

UniProt ID: Q8CIG8

UniProt Link: www.uniprot.org/uniprotkb/Q8CIG8/entry

Species Source: Mouse

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 637 AA

Molecular Weight: 73 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MAAMAVGGAGGSRVSSGRDLNCVPEIADTLGAVAKQGDFLCMPVFHPRFKREFIQEPAKNRPGP
QTRSDLLLSGRDWNTLIVGKLSPWIHPDSKVEKIRRNSEAAMLQELNFGAYLGLPAFLLPLNQEDNT
NLARVLTNHIHTGHHSSMFWMRVPLVAPEDLRDDVIANAPTTHTEEYSGEEKTWMWWHNFRTLCD
YSKRIAAVALEIGADLPSNHVIDRWLGEPIKAAILPTSIFLTNKKGFVLSKVQQRLIFRLLKLEVQFIITGT
NHHSEKEFCSYLQYLEYLSQNRPPPNAYELFAKGYEDYLSPLQPLMDNLESQTYEVFEKDPIKYSQ
YQQAIYKCLLDRVPEEEKETNVQVLMVLGAGRGPLVNASLRAAKQAERRIRLYAVEKNPNAVVTLEN
WQFEEWGSQVTVVSSDMREWVAPEKADIIVSELLGSFADNELSPECLDGAQHFLKDDGVSIPGEYT
SFLAPISSSKLYNEVRACREKDRDPEAQFEMPYVVRHLNHFHQLSAPKPCFTFSHPNRDPMIDNNRY
CTLEFPVEVNTVLHGFAFYFETVLYRDITLSIRPETHSPGMFSWFPIFFPIKQPITVHEGQNICVRFWR
CSNSKKVWYEWAVTAPVCSSIHNPTRSYTIGL

3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.



Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant Protein arginine N-methyltransferase 5 (Prmt5) (EC 2.1.1.320) (Histone-arginine N-methyltransferase PRMT5) (Jak-binding protein 1) (Shk1 kinase-binding protein 1 homolog) (SKB1 homolog) produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: Arginine methyltransferase that can both catalyze the formation of omega-N monomethylarginine (MMA) and symmetrical dimethylarginine (sDMA), with a preference for the formation of MMA (PubMed:15485929, PubMed:19584108, PubMed:19858291, PubMed:21917714, PubMed:23133559, PubMed:28263986). Specifically mediates the symmetrical dimethylation of arginine residues in the small nuclear ribonucleoproteins Sm D1 (SNRPD1) and Sm D3 (SNRPD3); such methylation being required for the assembly and biogenesis of snRNP core particles. Methylates SUPT5H and may regulate its transcriptional elongation properties (By similarity). May methylate the N-terminal region of MBD2 (By similarity). Mono- and dimethylates arginine residues of myelin basic protein (MBP) in vitro. May play a role in cytokine-activated transduction pathways. Negatively regulates cyclin E1 promoter activity and cellular proliferation (By similarity). Methylates histone H2A and H4 'Arg-3' during germ cell development (PubMed:16699504). Methylates histone H3 'Arg-8', which may repress transcription (PubMed:15485929). Methylates the Piwi proteins (PIWIL1, PIWIL2 and PIWIL4), methylation of Piwi proteins being required for the interaction with Tudor domain-containing proteins and subsequent localization to the meiotic nuage (PubMed:19584108). Methylates RPS10 (By similarity). Attenuates EGF signaling through the MAPK1/MAPK3 pathway acting at 2 levels. First, monomethylates EGFR; this enhances EGFR 'Tyr-1197' phosphorylation and PTPN6 recruitment, eventually leading to reduced SOS1 phosphorylation. Second, methylates RAF1 and probably BRAF, hence destabilizing these 2 signaling proteins and reducing their catalytic activity (PubMed:21917714). Required for induction of E-selectin and VCAM-1, on the endothelial cells surface at sites of inflammation. Methylates HOXA9. Methylates and regulates SRGAP2 which is involved in cell migration and differentiation (By similarity). Acts as a transcriptional corepressor in CRY1-mediated repression of the core circadian component PER1 by regulating the H4R3 dimethylation at the PER1 promoter (PubMed:23133559). Methylates GM130/GOLGA2, regulating Golgi ribbon formation. Methylates H4R3 in genes involved in glioblastomagenesis in a CHTOP- and/or TET1-dependent manner. Symmetrically methylates POLR2A, a modification that allows the recruitment to POLR2A of proteins including SMN1/SMN2 and SETX. This is required for resolving RNA-DNA hybrids created by RNA polymerase II, that form R-loop in transcription terminal regions, an important step in proper transcription termination. Along with LYAR, binds the promoter of gamma-globin HBG1/HBG2 and represses its expression. Symmetrically methylates NCL. Methylates p53/TP53; methylation might possibly affect p53/TP53 target gene specificity (By similarity). Involved in spliceosome maturation and mRNA splicing in prophase I spermatocytes through the catalysis of the symmetrical arginine dimethylation of SNRPB (small nuclear ribonucleoprotein-associated protein) and the interaction with tudor domain-containing protein TDRD6 (PubMed:28263986). Acts as a negative regulator of cGAS-STING signaling by mediating methylation of CGAS, preventing DNA-binding ability of CGAS (By similarity).

Subcellular Localization: Cytoplasm . Nucleus . Golgi apparatus . Note=Localizes to promoter regions of target genes on chromosomes (By similarity). Localizes to methylated chromatin (By similarity).

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer- composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.