

Recombinant Rat M3K8 Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP52501

Size: 10ug, 100ug

Protein Name: Mitogen-activated protein kinase kinase kinase 8 (EC 2.7.11.25) (Tumor progression locus 2) (TPL-2)

Synonym: Map3k8, Tpl2

UniProt ID: Q63562

UniProt Link: www.uniprot.org/uniprotkb/Q63562/entry

Species Source: Rat

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 467 AA

Molecular Weight: 53 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MEYMSTGSDEKEEIDLLINHLNVSEVLDIMENLYASEEPAVYEPSLMTMCPDSNQNKEHSESLLRSG
QEV PWLSSVRYGTVEDLLAFANHISNTTKHFYRCRPQESGILLNMVISPNQNGRYQIDSDVLLVPWKL
TYRSIGSGFVPRGAFGKVYLAQDMKTKKRMACKLIPVDQFKPSDVEIQACFRHENIAELYGAVLWGD
TVHLFMEAGEGGSVLEKLESCGPMREFEIIWVTKHVLKGLDFLHSSKKVIHHDIKPSNIVFMSTKAVLV
DFGLSVQMTEDVYLPKDLRGTEIYMSPEVILCRGHSTKADIYSLGATLIHMQTGTPPWVKRYPR SAY
PSYLYIIHKQAPPLEDIAGDCSPGMRELIEAALERNPNHRPKAADLLKHEALNPPREDQPRCQSLDSA
LFDRKRLLSRKELELPENIADSSCTGSTEESEVLRQRSLYIDLGALAGYFNIVRGPPTLEYG

3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.

Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant Mitogen-activated protein kinase kinase kinase 8 (EC 2.7.11.25) (Tumor progression locus 2) (TPL-2) produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: Required for lipopolysaccharide (LPS)-induced, TLR4-mediated activation of the MAPK/ERK pathway in macrophages, thus being critical for production of the pro-inflammatory cytokine TNF (TNF) during immune responses. Involved in the regulation of T-helper cell differentiation and IFNG expression in T-cells. Involved in mediating host resistance to bacterial infection through negative regulation of type I interferon (IFN) production. Transduces CD40 and TNFRSF1A signals that activate ERK in B-cells and macrophages, and thus may play a role in the regulation of immunoglobulin production. May also play a role in the transduction of TNF signals that activate JNK and NF-kappa-B in some cell types. In adipocytes, activates MAPK/ERK pathway in an IKBKB-dependent manner in response to IL1B and TNF, but not insulin, leading to induction of lipolysis. Plays a role in the cell cycle.

Subcellular Localization: Cytoplasm .

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer-composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.