

Recombinant Mouse EDA Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP51168

Size: 10ug, 100ug

Protein Name: Ectodysplasin-A (EDA protein homolog) (Tabby protein) [Cleaved into: Ectodysplasin-A, membrane form; Ectodysplasin-A, secreted form]

Synonym: Eda, Ed1, Ta

UniProt ID: O54693

UniProt Link: www.uniprot.org/uniprotkb/O54693/entry

Species Source: Mouse

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 391 AA

Molecular Weight: 42 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MGYPEVERREPLPAAAPRERGSQGCGRGAPARAGEGNSCRLFLGFFGLSLALHLLTLCCYLELRS
ELRRERGTESRLGGPGAPGTSGTLSSPGSLDPVGPITRHLGQPSFQQQPLEPGEDPLPPDSQDRH
QMA LLNFFFPDEKAYSEEESSRRVRNRKRSKSGEGADGPVKNKKKGKKAGPPGPNGPPGPPGPPG
PQGPPGIPGIPGIPGTTVMGPPGPPGPPGPPGPPGLQGPPSGAADKTGTRENQPAVHLQGQGS
QVKN DLSSGVLNDWSRITMNPVKVFLHPRSGEVLVDGTYFIYSQVEVYYINFTDFASYEVVVDEK
PFLQCTRSIETGKTNYNTCYTAGVCLLKARQKIAVKMVHADISINMSKHTTFFGAIRLGEAPAS

3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.

Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant Ectodysplasin-A (EDA protein homolog) (Tabby protein) [Cleaved into: Ectodysplasin-A, membrane form; Ectodysplasin-A, secreted form] produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: Cytokine which is involved in epithelial-mesenchymal signaling during morphogenesis of ectodermal organs. Functions as a ligand activating the DEATH-domain containing receptors EDAR and EDA2R. Isoform TAA binds only to the receptor EDAR, while isoform TA-A2 binds exclusively to the receptor EDA2R (By similarity). May also play a role in cell adhesion (PubMed:10534613).; Isoform TAA binds only to the receptor EDAR, while isoform TA-A2 binds exclusively to the receptor EDA2R.; Isoform TA-A2 binds exclusively to the receptor EDA2R.

Subcellular Localization: Cell membrane ; Single-pass type II membrane protein .; [Ectodysplasin-A, secreted form]: Secreted .

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer-composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.