

Recombinant Human ITB2 Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP50577

Size: 10ug, 100ug

Protein Name: Integrin beta-2 (Cell surface adhesion glycoproteins LFA-1/CR3/p150,95 subunit beta) (Complement receptor C3 subunit beta) (CD antigen CD18)

Synonym: ITGB2, CD18, MFI7

UniProt ID: P05107

UniProt Link: www.uniprot.org/uniprotkb/P05107/entry

Species Source: Human

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 769 AA

Molecular Weight: 85 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MLGLRPPLLALVGLLSLGCVLSECTKFKVSSCRECIESGPGCTWCQKLNFTGPGDPDSIRCDTRP
QLLMRGCAADDIMDPTSLAETQEDHNGGQKQLSPQKVTLYLPGQAAAFNVTFERRAKGYPIDLYYL
MDLSYSMLDDLNRNVKKLGDDLRLALNEITESGRIGFGSFVDKTVLPFVNTHPDKLRNPCPNKEKECQ
PPFAFRHVLKLTNNSNQFQTEVGKQLISGNLDAPEGGLDAMMQVAACPPEEIGWRNVTRLLVFATDD
GFHFAGDGKLGAILTPNDGRCHLEDNLYKRSNEFDYPSVGQLAHKLAENNIQPIFAVTSRMVKTYEK
LTEIIPKSAVGELSESSNVVHLIKNAYNKLSSRVFLDHNALPDTLKVTYDSFCSNGVTHRNQPRGDC
DGVQINVPITFQVKVTATECIEQSFVIRALGFTDIVTVQVLPQCECRCRDQSRDRSLCHGKGFLECG
ICRCDTGYIGKNCECQTQGRSSQELEGSCRKDNNSIICSGLGDCVCGQCLCHTSDVPGKLIYGQYC
ECDTINCERYNGQVCGGPGRGLCFCKGCRCHPGFEGSACQCERTTEGCLNPRRVECSGRGRRCRC
NVCECHSGYQLPLCQECPCSPCGKYISCAECLKFEKGPFCKNCSAACPLQLSNNPVKGRCTCK
ERDSEGCWVAYTLEQQDGM DRYLIYVDESRECVAGPNIAAIVGGTVAGIVLIGILLLVWKAHLHSDL
REYRRFEKEKLKSQWNNDNPLFKSATTVMNPKFAES

3. Buffer Composition



Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.

Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant Integrin beta-2 (Cell surface adhesion glycoproteins LFA-1/CR3/p150,95 subunit beta) (Complement receptor C3 subunit beta) (CD antigen CD18) produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: Integrin ITGAL:ITGB2 is a receptor for ICAM1, ICAM2 and ICAM3 (PubMed:1676048, PubMed:23775590, PubMed:38195629). Integrin ITGAL:ITGB2 is also a receptor for the secreted form of ubiquitin-like protein ISG15; the interaction is mediated by ITGAL (PubMed:29100055). Integrins ITGAM:ITGB2 and ITGAX:ITGB2 are receptors for the iC3b fragment of the third complement component and for fibrinogen. Integrin ITGAX:ITGB2 recognizes the sequence G-P-R in fibrinogen alpha-chain. Integrin ITGAM:ITGB2 recognizes P1 and P2 peptides of fibrinogen gamma chain. Integrin ITGAM:ITGB2 is also a receptor for factor X. Integrin ITGAD:ITGB2 is a receptor for ICAM3 and VCAM1 (PubMed:10438935, PubMed:8777714, PubMed:9841932). Contributes to natural killer cell cytotoxicity (PubMed:15356110). Involved in leukocyte adhesion and transmigration of leukocytes including T-cells and neutrophils (PubMed:11812992, PubMed:28807980). Triggers neutrophil transmigration during lung injury through PTK2B/PYK2-mediated activation (PubMed:18587400). Integrin ITGAL:ITGB2 in association with ICAM3, contributes to apoptotic neutrophil phagocytosis by macrophages (PubMed:23775590). In association with alpha subunit ITGAM/CD11b, required for CD177-PRTN3-mediated activation of TNF primed neutrophils (PubMed:21193407). Integrins ITGAX:ITGB2 functions as a receptor of the erythrocyte-specific adhesion molecule ICAM4 and mediates erythrophagocytosis (PubMed:16985175). Integrins ITGAX:ITGB2 functions as a receptor of the neuron-specific adhesion molecule ICAM5 ensuring neuron cell-leukocyte adhesion (PubMed:10741396). Integrin ITGAL:ITGB2 functions as a receptor of ICAM1 by acting as a platform at the immunological synapse to translate TCR engagement and density of the ITGAL ligand ICAM1 into graded adhesion (PubMed:38195629). Integrin ITGAM:ITGB2/MAC-1 complex functions as a signaling receptor for the ligand receptor ICAM1, ensuring adhesion between stimulated neutrophils and stimulated endothelial cells (PubMed:1980124). Integrin ITGAL/ITGB2 that functions as a signaling receptor of ICAM2, ensuring leukocyte cell-cell adhesion on resting cells (PubMed:1676048).

Subcellular Localization: Cell membrane ; Single-pass type I membrane protein . Membrane raft ; Single-pass type I membrane protein .

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer-composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial

manufacturing.