

Recombinant Rat HNRDL Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP52695

Size: 10ug, 100ug

Protein Name: Heterogeneous nuclear ribonucleoprotein D-like (hnRNP D-like) (hnRNP DL)

Synonym: Hnrnpdl, HnrpdI

UniProt ID: Q3SWU3

UniProt Link: www.uniprot.org/uniprotkb/Q3SWU3/entry

Species Source: Rat

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 322 AA

Molecular Weight: 35 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MTGTARSALPLPQSPARALRPSGAARAAPSLSPSRFSACPLDPSSFPTSGNKMFIGGLSWDTSKDD
LTEYLSRFGEVVDCTIKTDPVTGRSRGFGFVLFKDAASVDKVLKEHKLDGKLIDPKRAKALKGKEP
PKKV FVGG LSPDTSEEQIKEYFGAFGEIENIELPMDTKTNERRGFCFITYTDEEPPVKLLLESRYHQIGS
GKCEIKVAQPKEVYRQQQQQKGGRGAAAGGRGGARGRGQQGNWNQGFNNYYDQGYGNYN
SAYGDESYSGYGGYDYGYNYSYGYGGYTDYSGQQSTYKGASRGGGNHQNNYQPY

3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.

Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant Heterogeneous nuclear ribonucleoprotein D-like (hnRNP D-like) (hnRNP DL) produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: Acts as a transcriptional regulator. Promotes transcription repression. Promotes transcription activation in differentiated myotubes. Binds to double- and single-stranded DNA sequences. Binds to the transcription suppressor CATR sequence of the COX5B promoter. Binds with high affinity to RNA molecules that contain AU-rich elements (AREs) found within the 3'-UTR of many proto-oncogenes and cytokine mRNAs. Binds both to nuclear and cytoplasmic poly(A) mRNAs. Binds to poly(G) and poly(A), but not to poly(U) or poly(C) RNA homopolymers. Binds to the 5'-ACUAGC-3' RNA consensus sequence (By similarity).

Subcellular Localization: Nucleus . Cytoplasm . Note=Shuttles between the nucleus and the cytoplasm in a TNPO1-dependent manner.

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer-composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.