

# Recombinant Mouse CASP7 Protein

**For Research Use Only**

## 1. Basic Product Information

Catalog No.: REP51530

Size: 10ug, 100ug

Protein Name: Caspase-7 (CASP-7) (EC 3.4.22.60) (Apoptotic protease Mch-3) (Cysteine protease LICE2) [Cleaved into: Caspase-7 subunit p20; Caspase-7 subunit p11]

Synonym: Casp7, Lice2, Mch3

UniProt ID: P97864

UniProt Link: [www.uniprot.org/uniprotkb/P97864/entry](http://www.uniprot.org/uniprotkb/P97864/entry)

Species Source: Mouse

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 303 AA

Molecular Weight: 34 KD

Protein Tag: C-terminal His6

## 2. Amino Acid Sequence

MTDDQDCAAELEKVDSSSEDGVDAKPDRSSISSLKKKRNASAGPVRTGRDRVPTYLYRMDFQK  
MGKCIINNNKFDKATGMDVRNGTDKDGALFKCFQNLGFEVTVHNDSCAKMQDLLRKASEEDHS  
NSACFACVLLSHGEEDLIYGKDGVTPIKDLTAHFRGDRCKTLLEKPKLFFIQACRGTELDGDIQADSG  
PINDIDANPRNKIPVEADFLFAYSTVPGYYSWRNPNGKGSWFVQALCSILNEHGKDLEIMQILTRVNDR  
VARHFESQSDDPRFNEKKQIPCMVSMMLTKELYFSR

## 3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.

Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

## 4. Storage Conditions



Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

## 5. Product Description

This product is recombinant Caspase-7 (CASP-7) (EC 3.4.22.60) (Apoptotic protease Mch-3) (Cysteine protease LICE2) [Cleaved into: Caspase-7 subunit p20; Caspase-7 subunit p11] produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

**Protein Function:** Thiol protease involved in different programmed cell death processes, such as apoptosis, pyroptosis or granzyme-mediated programmed cell death, by proteolytically cleaving target proteins (PubMed:16469926, PubMed:19168786). Has a marked preference for Asp-Glu-Val-Asp (DEVD) consensus sequences, with some plasticity for alternate non-canonical sequences (PubMed:25231987). Its involvement in the different programmed cell death processes is probably determined by upstream proteases that activate CASP7 (PubMed:18667412, PubMed:35705808). Acts as an effector caspase involved in the execution phase of apoptosis: following cleavage and activation by initiator caspases (CASP8 and/or CASP9), mediates execution of apoptosis by catalyzing cleavage of proteins, such as CLSPN, PARP1, PTGES3 and YY1 (PubMed:16469926). Compared to CASP3, acts as a minor executioner caspase and cleaves a limited set of target proteins (PubMed:16469926). Acts as a key regulator of the inflammatory response in response to bacterial infection by catalyzing cleavage and activation of the sphingomyelin phosphodiesterase SMPD1 in the extracellular milieu, thereby promoting membrane repair (PubMed:35705808). Regulates pyroptosis in intestinal epithelial cells: cleaved and activated by CASP1 in response to *S.typhimurium* infection, promoting its secretion to the extracellular milieu, where it catalyzes activation of SMPD1, generating ceramides that repair membranes and counteract the action of gasdermin-D (GSDMD) pores (PubMed:22807671, PubMed:35705808). Regulates granzyme-mediated programmed cell death in hepatocytes: cleaved and activated by granzyme B (GZMB) in response to bacterial infection, promoting its secretion to the extracellular milieu, where it catalyzes activation of SMPD1, generating ceramides that repair membranes and counteract the action of perforin (PRF1) pores (PubMed:35705808). Following cleavage by CASP1 in response to inflammasome activation, catalyzes processing and inactivation of PARP1, alleviating the transcription repressor activity of PARP1 (PubMed:22464733). Acts as an inhibitor of type I interferon production during virus-induced apoptosis by mediating cleavage of antiviral proteins CGAS, IRF3 and MAVS, thereby preventing cytokine overproduction (PubMed:30878284). Cleaves and activates sterol regulatory element binding proteins (SREBPs) (By similarity). Cleaves phospholipid scramblase proteins XKR4, XKR8 and XKR9 (PubMed:25231987). Cleaves BIRC6 following inhibition of BIRC6-caspase binding by DIABLO/SMAC (By similarity).

**Subcellular Localization:** Cytoplasm, cytosol . Nucleus . Secreted, extracellular space .

Note=Following cleavage and activation by CASP1 or granzyme B (GZMB), secreted into the extracellular milieu by passing through the gasdermin-D (GSDMD) pores or perforin (PRF1) pore, respectively.

## 6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer-composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.