

Recombinant Mouse KS6A5 Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP51751

Size: 10ug, 100ug

Protein Name: Ribosomal protein S6 kinase alpha-5 (S6K-alpha-5) (EC 2.7.11.1) (90 kDa ribosomal protein S6 kinase 5) (Nuclear mitogen- and stress-activated protein kinase 1) (RSK-like protein kinase) (RLSK)

Synonym: Rps6ka5, Msk1

UniProt ID: Q8C050

UniProt Link: www.uniprot.org/uniprotkb/Q8C050/entry

Species Source: Mouse

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 863 AA

Molecular Weight: 97 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MEGEGGGSGGAGTSGDSGDGGEQLLTVKHELRTANLTGHAEKVGIENFELLKVLGTGAYGKVFLV
RKISGHDAGKLYAMKVLKKATIVQKAKTTEHTRTERQVLEHIRQSPFLVTLHYAFQTETKLHLILDYIN
GGELFTHLSQRERFTEHEVQIYVGEIVLALEHLHLKLGIIYRDIKLENILLDSNGHVLTDFGLSKEFVAD
ETERAYSFCGTIYMAPDIVRGGDSGHDKAVDWWSLGVLMYELLTGASPFTVDGEKNSQAEISRIL
KSEPPYPQEMSTVAKDLLQRLLMKDPKKRLGCGPRDAEEIKEHLFFEKIKWDDLAACKVPAPFKPVI
RDELDVSNFAEEFTEMDPTYSPAALPQSSERLFQGYSFVAPSILFKRNAVIDPLQFHMGVDRPGVT
NVARSAMMKDSPFYQHYDLDLKDKPLGEGSFSICRKC VHKKTNQAFVAVKIISKRMEANTQKEITALKL
CEGHPNIVKLHEVFHDQVAASAQPPGQVVLCSLLLLALLFNRSLTRKPVTWTWLVHSTS QLPPLPPP
MPEIVLFILLSDNGQLHTFLVMELLNGGELFERIKRKKHFS ETEASYIMRKLVS AVSHMHDVG VVHRD
LKPENLLFTDENDNLEIKVIDFGFARLKPPDNQPLKTPCFTLHYAAPELLTHNGYDESCDLWSLGVILY
TMLSGQVPFQSHDRSLTCTSAVEIMKKIKKGDFSFEGEAWKNVSQEAKDLIQGLLTVDPNKRLKMSG
LRYNEWLQDGSQSSNPLMTPDILGSSGAHVHTCVKATFHAFNKYKREGFCLQNVDKAPLAKRRKM
KRTSTSTETRSSSSSESSRSSSSSQSHGKTTPTKTLQPSNPTEGSPDPTLFQFSD

3. Buffer Composition



Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.

Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant Ribosomal protein S6 kinase alpha-5 (S6K-alpha-5) (EC 2.7.11.1) (90 kDa ribosomal protein S6 kinase 5) (Nuclear mitogen- and stress-activated protein kinase 1) (RSK-like protein kinase) (RLSK) produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: Serine/threonine-protein kinase that is required for the mitogen or stress-induced phosphorylation of the transcription factors CREB1 and ATF1 and for the regulation of the transcription factors RELA, STAT3 and ETV1/ER81, and that contributes to gene activation by histone phosphorylation and functions in the regulation of inflammatory genes (By similarity) (PubMed:11553624, PubMed:11909979, PubMed:16806820). Phosphorylates CREB1 and ATF1 in response to mitogenic or stress stimuli such as UV-C irradiation, epidermal growth factor (EGF) and anisomycin (PubMed:11909979). Plays an essential role in the control of RELA transcriptional activity in response to TNF and upon glucocorticoid, associates in the cytoplasm with the glucocorticoid receptor NR3C1 and contributes to RELA inhibition and repression of inflammatory gene expression (PubMed:12628924, PubMed:16806820). In skeletal myoblasts is required for phosphorylation of RELA at 'Ser-276' during oxidative stress (PubMed:12628924). In erythropoietin-stimulated cells, is necessary for the 'Ser-727' phosphorylation of STAT3 and regulation of its transcriptional potential (PubMed:11553624). Phosphorylates ETV1/ER81 at 'Ser-191' and 'Ser-216', and thereby regulates its ability to stimulate transcription, which may be important during development and breast tumor formation (By similarity). Directly represses transcription via phosphorylation of 'Ser-1' of histone H2A (By similarity). Phosphorylates 'Ser-10' of histone H3 in response to mitogenics, stress stimuli and EGF, which results in the transcriptional activation of several immediate early genes, including proto-oncogenes c-fos/FOS and c-jun/JUN (PubMed:15870105, PubMed:16517600). May also phosphorylate 'Ser-28' of histone H3 (PubMed:11441012, PubMed:15870105). Mediates the mitogen- and stress-induced phosphorylation of high mobility group protein 1 (HMGN1/HMG14) (By similarity). In lipopolysaccharide-stimulated primary macrophages, acts downstream of the Toll-like receptor TLR4 to limit the production of pro-inflammatory cytokines (PubMed:18690222). Functions probably by inducing transcription of the MAP kinase phosphatase DUSP1 and the anti-inflammatory cytokine interleukin 10 (IL10), via CREB1 and ATF1 transcription factors (PubMed:18690222). Plays a role in neuronal cell death by mediating the downstream effects of excitotoxic injury (PubMed:12807421). Phosphorylates TRIM7 at 'Ser-106' in response to growth factor signaling via the MEK/ERK pathway, thereby stimulating its ubiquitin ligase activity (By similarity).

Subcellular Localization: Nucleus .

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer-composition requirements at the time of order placement.



Enlibio Wuhan Enlibio Biotech Co.,Ltd

- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.