

Recombinant Mouse BMP2 Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP51316

Size: 10ug, 100ug

Protein Name: Bone morphogenetic protein 2 (BMP-2) (Bone morphogenetic protein 2A) (BMP-2A)

Synonym: Bmp2, Bmp-2

UniProt ID: P21274

UniProt Link: www.uniprot.org/uniprotkb/P21274/entry

Species Source: Mouse

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 394 AA

Molecular Weight: 45 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MVAGTRCLLVLLL PQVLLGGAAGLIPELGRKKFAAASSRPLSRPSEDVLSEFELRLLSMFGLKQRPTP
SKDVVVPYMLDLYRRHSGQPGAPAPDHRLEAASRANTVRSFHHEEAVEELPEMSGKTARRFFF
NLSSVPSDEFLTSAELQIFREQIQEALGNSSFQHRINIYEIIPAAANLKFVTRLLDTRLVNQNTSQWE
SFDVTPAVMRWTTQGHTNHGFVVEVAHLEENPGVSKRHVRISRSLHQDEHSWSQIRPLLVTFGHD
GKGHPLHKREKRQAKHKQRKRLKSSCKRHPLYVDFSDVGWNDWIVAPPGYHAFYCHGECPPFLAD
HLNSTNHAIVQTLVNSVNSKIPKACCVPTELSAISMLYLDENEKVVLKNYQDMVVEGCGCR

3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.

Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant Bone morphogenetic protein 2 (BMP-2) (Bone morphogenetic protein 2A) (BMP-2A) produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: Growth factor of the TGF-beta superfamily that plays essential roles in many developmental processes, including cardiogenesis, neurogenesis, and osteogenesis. Induces cartilage and bone formation. Initiates the canonical BMP signaling cascade by associating with type I receptor BMPR1A and type II receptor BMPR2. Once all three components are bound together in a complex at the cell surface, BMPR2 phosphorylates and activates BMPR1A. In turn, BMPR1A propagates signal by phosphorylating SMAD1/5/8 that travel to the nucleus and act as activators and repressors of transcription of target genes. Also acts to promote expression of HAMP, via the interaction with its receptor BMPR1A/ALK3 (PubMed:31800957). Can also signal through non-canonical pathways such as ERK/MAP kinase signaling cascade that regulates osteoblast differentiation. Also stimulates the differentiation of myoblasts into osteoblasts via the EIF2AK3-EIF2A-ATF4 pathway by stimulating EIF2A phosphorylation which leads to increased expression of ATF4 which plays a central role in osteoblast differentiation. Acts as a positive regulator of odontoblast differentiation during mesenchymal tooth germ formation, expression is repressed during the bell stage by MSX1-mediated inhibition of CTNNB1 signaling (PubMed:29148101).

Subcellular Localization: Secreted.

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer-composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.