

Recombinant Rat MK09 Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP52374

Size: 10ug, 100ug

Protein Name: Mitogen-activated protein kinase 9 (MAP kinase 9) (MAPK 9) (EC 2.7.11.24) (SAPK-alpha) (Stress-activated protein kinase JNK2) (c-Jun N-terminal kinase 2) (p54-alpha)

Synonym: Mapk9, Jnk2, Prkm9

UniProt ID: P49186

UniProt Link: www.uniprot.org/uniprotkb/P49186/entry

Species Source: Rat

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 423 AA

Molecular Weight: 48 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MSDSKSDGQFYSVQVADSTFTVLKRYQQLKPIGSGAQQGIVCAAFDTVLGINVAVKKLSRPFQNQTHA
KRAYRELVLLKCVNHKNIISLLNVFTPQKTLEEFQDVYLVMEMLDANLCQVIHMELDHERMSYLLYQM
LCGIKHLHSAGIIHRDLKPSNIVKSDCTLKILDFGLARTACTNFMMPYVVTRYRAPEVILGMGYKE
NVDIWSVGCIMGELVKGCVIFQGTDHIDQWNKVIEQLGTPSAEFMKKLQPTVRNYVENRPKYPGIKF
EELFPDWIFPSESERDKIKTSQARDLLSKMLVIDPDKRISVDEALRHPYITVWYDPAEAEAPPPQIYDA
QLEEREHAIEEWKELIYKEVMDWEERSKNGVKDQPSDAAVSSKATPSQSSSINDISSMSTEHTLASD
TDSSLDASTGPLEGCR

3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.

Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant Mitogen-activated protein kinase 9 (MAP kinase 9) (MAPK 9) (EC 2.7.11.24) (SAPK-alpha) (Stress-activated protein kinase JNK2) (c-Jun N-terminal kinase 2) (p54-alpha) produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: Serine/threonine-protein kinase involved in various processes such as cell proliferation, differentiation, migration, transformation and programmed cell death. Extracellular stimuli such as pro-inflammatory cytokines or physical stress stimulate the stress-activated protein kinase/c-Jun N-terminal kinase (SAP/JNK) signaling pathway. In this cascade, two dual specificity kinases MAP2K4/MKK4 and MAP2K7/MKK7 phosphorylate and activate MAPK9/JNK2. In turn, MAPK9/JNK2 phosphorylates a number of transcription factors, primarily components of AP-1 such as JUN and ATF2 and thus regulates AP-1 transcriptional activity. In response to oxidative or ribotoxic stresses, inhibits rRNA synthesis by phosphorylating and inactivating the RNA polymerase 1-specific transcription initiation factor RRN3. Promotes stressed cell apoptosis by phosphorylating key regulatory factors including TP53 and YAP1. In T-cells, MAPK8 and MAPK9 are required for polarized differentiation of T-helper cells into Th1 cells. Upon T-cell receptor (TCR) stimulation, is activated by CARMA1, BCL10, MAP2K7 and MAP3K7/TAK1 to regulate JUN protein levels. Plays an important role in the osmotic stress-induced epithelial tight-junctions disruption. When activated, promotes beta-catenin/CTNNB1 degradation and inhibits the canonical Wnt signaling pathway. Also participates in neurite growth in spiral ganglion neurons. Phosphorylates the CLOCK-BMAL1 heterodimer and plays a role in the regulation of the circadian clock (By similarity). Phosphorylates POU5F1, which results in the inhibition of POU5F1's transcriptional activity and enhances its proteasomal degradation (By similarity). Phosphorylates ALKBH5 in response to reactive oxygen species (ROS), promoting ALKBH5 sumoylation and inactivation (By similarity).

Subcellular Localization: Cytoplasm . Nucleus . Note=Colocalizes with POU5F1 in the nucleus.

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer-composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.